

Assessment of Maldivian Coral Reefs in 2009 after Natural Disasters



Hussein Zahir
Norman Quinn
Nicky Cargillia



Marine Research Centre
Ministry of Fisheries and Agriculture
Malé, Republic of Maldives

Assessment of Maldivian Coral Reefs in 2009 after Natural Disasters

Hussein Zahir, Norman Quinn and Nicky Cargillia

**Marine Research Centre
Ministry of Fisheries and Agriculture, Malé, Republic of Maldives**

Copy right© 2010 Marine Research Centre. Ministry of Fisheries and Agriculture, Malé, Republic of Maldives

Citations: Zahir, H¹; N. Quinn² and N. Cargilia³ (2010). Assessment of Maldivian Coral Reefs in 2009 after Natural Disasters. Marine Research Centre; Malé; Republic of Maldives. v + 57 pp.

¹Senior Reef Ecologist, Marine Reseach Centre, Ministry of Fisheries and Agriculture. Male, Republic of Maldives; ²Fulbright Scholar at Marine Research Centre from 2008-2009, ³Intern at MRC in 2009 under Maldives Environmental Management Project (MEMP) of Maldives

Available from: Marine Research Centre

Postal Address: Marine Research Centre,
H. White Waves, Moonlight Higung, 20025,
Malé, Republic of Maldives

Telephone: + (960) 332 2242
Facsimile: + (960) 332 2509
Email: info@mrc.gov.mv
Internet: www.mrc.gov.mv

Cover Photo: Marine Research Center staff members M. Shafiya and H. Hamid observing a school of Oriental Sweetlips (*Plectorhinchus vittatus*) on a training dive at 15 m along Vadoo Canyons, South Male' atoll, January 2009.

Table of Contents

List of Abbreviations	1
Foreword	2
Acknowledgements	3
Report Objectives	4
Introduction	4
Initial Reef Check Surveys and the 1998 Bleaching Event	5
Tsunami Devastation	7
Erosion	6
Natural Resilience - Recruitment	8
National Coral Reef Monitoring Survey Sites	9
Survey Methods	12
Results	13
Reef recovery: Coral cover	13
Haa Dhaalu Atoll	13
North and South Male' Atoll	16
<i>Maniyafushi</i>	16
<i>En'boodhoofinolhu</i>	16
<i>Bodu Ban'dos</i>	19
<i>Udhafushi</i>	19
Depth Variation	19
Ari Atoll	20
<i>Fesdhoo, Maayafushi, Velidhoo reefs</i>	20
<i>Dega Giri</i>	21
<i>Kandholhudhoo</i>	21
Vaavu Atoll	22
Gaafu Alifu Atoll	24
Addu Atoll	26
Coral Framework Status	28
Coral Reef Recovery in the Region	30
Dubai, United Arab Emirates	30
Lakshadweep Islands, India	31
Chagos Archipelago, Indian Ocean	31
Coral Reef Diseases	32

Fish Distribution and Abundance	32
Grouper Abundance	34
Other Reef Fish	34
Sponge Abundance and Distribution	34
Echinoderms	35
Crown of Thorns starfish - <i>Acanthaster planci</i>	35
Echinoids	37
Holothurians Abundance and Distribution	37
Crustacean Abundance and Distribution	40
Lobsters	40
Banded coral shrimp	41
Mollusc - Abundance and Distribution	41
Giant Clams – <i>Tridacna spp.</i>	41
Triton Shells	43
Anthropogenic Disturbances	43
Sea Water Temperature Variation	44
Adaption to Sea Water Temperature Variation	46
Future Coral Reef Monitoring Efforts and Awareness Development	47
References	50

List of Figures

Figure 1	Comparison of NCRMS 150 m point intercept transect percent live coral cover data at three sites in 1997 before and immediately after the bleaching in April – May 1998 bleaching event (after Zahir 2002a)	5
Figure 2	A sea cucumber (<i>Holothurian atra</i>) lies among the rubble of the devastated reef flat (2 m) at Vabbinfaru reef, September 2009	6
Figure 3	Percent live coral cover reported by surveys undertaken in January – February 2005, a couple months after the December 2004, tsunami (after AusAID, 2005)	7
Figure 4	Trees fall into the sea as waves on Vaavu atoll sweep across the reef eroding Bodumohora Island, March 2009	8
Figure 5	During high tide with strong westerly wind the sea washes with several meters of the villas, North Male’ atoll, October 2009	8
Figure 6	(a-f) 6a) Map of Maldives with yellow colored areas showing areas with NCRMS sites sampled in 2009. 6b) Map of most northern NCRMS sites in Haa Dhaalu atoll, 6c) Map of central NCRMS sites in North and South Male’ atolls, 6d) Map of central NCRMS sites in Ari atoll, 6e) Map of central NCRMS sites in Vaavu atoll and 6f) Map of most southern NCRMS sites in Addu atoll. Orange indicates islands. Only site names are shown in each atoll for clarity, yellow circles indicate general location of the transects	10
Figure 7	Mean (3 m, 10 m) percent coral cover of the 17 NCRMS sites surveyed from March to October 2009. Error bars show standard deviation. n = 8 for each site	14

Figure 8	Percentage live coral cover at 3 m at three sites in Haa Dhaalu atoll. Error bars only included for 2009, n = 4 for each site	14
Figure 9	Comparison of % coral cover at 3 m and 10 m at three sites at Haa Dhaalu atoll, March 2009. n = 4. Standard error shown by bars	15
Figure 10	Percent cover of the corallimorpharian population <i>Discosoma c.f. rhodostoma</i> on Hirimaradhoo reef at 3 m from 1998 to 2009. Standard error bars only shown for 2009 (n = 4)	15
Figure 11	Schools of convict surgeon fish, <i>Acanthurus triosteges</i> , at 5 m, Hon'doofushi, Haa Dhaalu, March 2009	16
Figure 12	A boat channel was constructed through the Hirimaradhoo reef after the 2005 survey	16
Figure 13	Debris tossed from Maniyafushi jetty covers part of the reef, February 2009	17
Figure 14	Percent live coral cover at 3 m at five sites in North and South Male' atoll. Standard error bars only included for 2009. n = 4 for each site	17
Figure 15	M. A. Abdulla and B. L. Kojis swim past healthy <i>Acropora</i> colonies as they return from conducting a survey at Maniyafushi reef, February 2009	18
Figure 16	Recently killed juvenile <i>Acropora sp.</i> coral. Possibly killed by <i>Culcita schmideliana</i> or a coral eating gastropod	18
Figure 17	A swarm of coral eating snails have eaten an <i>Acropora sp.</i> colony	18
Figure 18	The pin cushion starfish, <i>Culcita schmideliana</i> , on reef. This starfish eats small corals	18
Figure 19	Damaged <i>Acropora</i> at En'boodhoofinolhu reef, South Male' atoll, at 10 m, September, 2009	20
Figure 20	Comparison of % coral cover at 3 m and 10 m at one site at North Male' atoll and two sites at South Male' atoll, in 2009. n = 4 at each site and each depth. Standard error shown by bars	20
Figure 21	Percentage live coral cover at five sites in Ari atoll. Standard error bars only included for 2009. (n = 4 for each site). Dega giri data from Solandt and Wood (2008)	21
Figure 22	Kandholhudhoo is a picnic island with beautiful beaches naturally replenished by a diverse, healthy coral reef which surrounds the island. April 2009	21
Figure 23	Kandholhudho reef at 7 m showing some physical damage and partial mortality to tabulate <i>Acropora</i> and intact tabulate <i>Acropora</i> colonies, March 2009	22
Figure 24	Comparison of % coral cover at 3 m and 10 m at four sites at Ari atoll, April 2009. n = 4 at each site and each depth. Standard error shown by bars	22
Figure 25	Percentage live coral cover at three sites in Vaavu Atoll. Standard error bars only included for 2009, n = 4 for each site and year. 1999 point represents pooled values from McClanahan (2000)	23
Figure 26	Comparison of percent coral cover at 3 m and 10 m at three sites at Vaavu atoll, May 2009. n = 4 at each site and depth. Standard error shown by bars	23
Figure 27	Percentage live coral cover at 5 m at Koodoo, and Han'dahaa in Gaafu Alifu atoll. n = 4 for Koodoo for each year. Han'dahaa based on visual estimate survey	24
Figure 28	Over lapping of tabulate corals (<i>Acropora cytherea</i>) at 10 m at Han'dahaa, Gaafu Alifu atoll has created one of the few sites known where coral cover exceeds 100%. September 2009	24
Figure 29	<i>Acropora</i> dominated the north east reef slope at 15 - 30 m, Han'dahaa, Gaafu Alifu atoll, September 2009	25
Figure 30	Dislodged and overturned <i>Acropora</i> corals on reef flat, 4 m, Han'dahaa reef, September 2009. Note corals are still alive and new horizontal growth is occurring on left coral	25
Figure 31	S. Ali inspects filamentous blue green algae on a tabulate coral and recently dead tabulate coral (white area) on the Han'dahaa reef flat near the desalinization plant outfall (see upper left hand corner of photo). 4 m, September 2009	25
Figure 32	Y. Rilwan is recording substrate cover on a transect line at Vilingili reef, Addu atoll. 10 m, May 2009	26

Figure 33	Percentage live coral cover at 5 m at Gan, Vilingili and Hithadhoo reef, Addu atoll. Standard error bars only included for 2009, n = 4 for each site and year	26
Figure 34	Recently dead coral at Vilingili reef, Addu atoll, May 2009. Algae are beginning to grow on portions of the dead coral	26
Figure 35	Large colonies of <i>Porites lobata</i> were common at Vilingili reef, May 2009	27
Figure 36	Percentage live coral cover at 10 m at three sites in Addu atoll, May 2009. n = 4 at each site in 2009 and n = 3 in 2002 – 2004	27
Figure 37	I. Abid, a member of the MRC coral reef monitoring team conducting a survey along four 20 m transects at Hithadhoo reef, Addu atoll in May 2009	27
Figure 38	Comparison of % coral cover at 3 m and 10 m at three sites at Addu atoll, May 2009. n = 4 at each site and each depth. Standard error shown by bars	28
Figure 39	Dead <i>Porites</i> colony with new colonies of <i>Porities</i> , <i>Acropora</i> and other species growing on it, 20 m, Magiri reef, North Male' atoll, September 2009	29
Figure 40	Guraidhoo Corner, South Male' atoll is a reef subject to oceanic swells and unconsolidated corals with poor recovery. September 2009, 8 m	30
Figure 41	Coralline algae being attacked by the bacterial disease, CLOD (yellow), at En'boodhoofinolhu, North Male' atoll, September 2009. White section is recently killed coralline algae	32
Figure 42	a-e Average abundance of each Reefcheck fish category at the NCRM survey sites in each atoll	33
Figure 43	Average abundance per category of fish in each atoll surveyed	34
Figure 44	Geographic and depth variation of % sponge cover in NCRMS sites in 2009	35
Figure 45	<i>Terpios sp.</i> overgrowing <i>Acropora</i> coral, Kandholhudhoo, Ari atoll, 10 m, April 2009	35
Figure 46	Dead juvenile Crown-of-Thorns starfish, <i>Acanthaster planci</i> , from Bodumohora reef, Vaavu Atoll, October 2008	36
Figure 47	Monthly records of <i>Acanthaster planci</i> and <i>Culcita schmideliana</i> collected from Vabbinfaru reef in 2006 and 2007 (Azeez, pers. comm.)	36
Figure 48	Monthly records of <i>Acanthaster planci</i> and <i>Culcita schmideliana</i> collected from Ihuru reef, North Male' atoll in 2006 and 2007 (Azeez, pers. comm.)	37
Figure 49	Abundance and distribution of two urchin species in the NCRMS sites in 2009	37
Figure 50	Total abundance of sea cucumber individuals spotted at each of the surveyed atolls including Reefcheck and free roving dives	38
Figure 51	<i>Thelenota ananas</i> at 20 m at Coral Gardens, North Male' atoll, October 2009	39
Figure 52	Post puerulis lobster on Gan reef, 5 m, May 2009. Carapace length estimated at 1 cm	40
Figure 53	Giant clam, <i>Tridacna squamosa</i> , about 35 cm long at Vattaru, Vaavu atoll, 10 m, March 2009	41
Figure 54	H. Zahir, photographing a large <i>Tridacna squamosa</i> (TL 45 cm) at Hithadhoo reef, 10 m, Addu atoll, May 2009	42
Figure 55	Abundance distribution of giant clams, <i>Tridacna spp.</i> , in NCRMS sites at shallow and deep transects in 2009	42
Figure 56	Size frequency distribution of giant clams, <i>Tridacna spp.</i> , from the 2009 NCRMS	43
Figure 57	Worker fumigating on island resort, North Male' atoll, September 2009	44
Figure 58	Mean satellite derived surface sea water temperature for the central atolls from 1950 – 1999 + 4 S.D. (after Edwards, 2001). Specific site data recorded between 10 - 25 m in 2009	45
Figure 59	Mean S3T at coral nursery (3 m) at Huvafenfushi, North Male' atoll, from August 2007 to February 2008	45
Figure 60	Mean monthly sea water temperatures at 8 m and 16 m from Rasdhoo, Ari atoll, April 2008 – March 2009 compared with Edwards, et al. (2001) SST	46
Figure 61	Temperature by depth profile at Barakuda Point, September 2009. Y axis on the top figure is depth in meters and the oC in the bottom figure. The x axis represents dive time in minutes 10 minutes from the start of dive at 9:40am	46
Figure 62	South Male' atoll, outer reef, 15 m, August 2009. Future surveys need to include coral communities on the outer reef slope of the atolls. Lateral visibility estimated at over 35 m	47

Figure 63 Photosynthesis is a process that converts carbon dioxide into organic compounds, especially sugars, using the energy from sunlight. In the sea, instead of a tomato plant, photosynthesis is done by plankton, coral, algae and sea grass 49

List Tables

Table 1	National coral reef monitoring survey sites with geographic coordinates and atoll and regional classification of the sites. sites sampled in 2009 indicated with x. thuvuru, anbaraa, and maduvvari were sampled from 1997 – 1999	12
Table 2	List of species encountered on ncrm surveys and during free rover dives at >20m with the commercial value for each species (adapted from FAO statistics, conand 1990) . x indicates species present in survey	38

Abbreviations

AVHRR	Advanced Very High Resolution Radiometer
CoT	Crown of Thorns starfish – <i>Acanthaster planci</i>
GBR	Great Barrier Reef
IUCN	International Union for Conservation of Nature
MMS3T	Mean monthly sub surface temperature
MMSST	Mean monthly sea water temperature
MOFAMR	Ministry of Fisheries, Agriculture and Marine Resources
MPA	Marine Protected Areas
MRC	Marine Research Centre
NCRMS	National Coral Reef Monitoring Survey
SST	surface sea water temperature
S3T	Subsurface sea water temperature

The place names spellings used in the report are based on the Ministry of Planning and National Development's Official Atlas of the Maldives (2008)

Foreword

Coral reefs and its ecosystem services, from coastal protection through its physical structure, to fisheries through exploitation of multitude of marine fishes and tourism are of immense importance to Maldives for its physical integrity and economy. Maldivians have been living in this atoll archipelago for millennia, their culture and livelihood driven by the resources provided by the reefs and the surrounding ocean. These resources are now threatened by both unmanaged exploitation and climate change impacts.

Climate change related impacts are among one of the most concerning threats to the coral reefs. Among the major threats of climate change are coral bleaching, coral diseases and ocean acidification related impacts. In 1998 an intense El Niño increased sea surface temperatures of much of the western Pacific and Indian Oceans causing widespread coral bleaching leading to mass coral mortality in many countries. For example, in Palau, more than 90% of the corals on some reefs bleached and at least 50% perished. In the Maldives, bleaching caused coral cover in the shallow waters to plummet to only about 3%.

In addition to the global and regional effects of climate change, there are several localized threats and impacts to coral reef ecosystems. Reef fishes are intensely harvested, greatly reducing their abundances and altered food web dynamics on all but the most isolated or intensely managed reefs around the world. Additionally, some destructive fishing practices directly kill corals and can destroy the reef matrix reducing its resilience and making them vulnerable to additional stress. Reefs are also threatened by many other local activities such as coastal development that increase sediment runoff into coastal areas and coastal reefs, in some cases smothering and killing corals. Some scientists believe that these and other local impacts could act synergistically with climate change thereby magnifying the negative effects of human-induced and climate stressors.

The bleaching event of 1998 has led to the current impetus in systematic effort on coral reef monitoring by the Marine Research Centre. Monitoring of coral reefs at selected sites has been ongoing since 1998 to provide an understanding of reef recovery and factors influencing recovery following the bleaching event. This report is a consolidated effort of past 10 years of reef monitoring efforts. Despite the documented and projected impacts of climate change on coral reefs, there is reason to be optimistic. The data indicate that the potential for recovery is high despite the regional variability. First, it is possible that corals will acclimate or evolve to become more tolerant of rising temperature. Given the unprecedented rate of climate change and the suite of changes and stresses the corals would have to adapt to, many coral biologists are skeptical that adaptation and acclimation alone will facilitate long-term survival if climate related threats are not abated.

In this context the existing Maldives coral reef monitoring program needs to be expanded to cover more reefs and allow detailed examination of the impacts of climate change on coral reefs and associated ecosystems currently recovering after the massive bleaching event of 1998. To facilitate increased monitoring of reef resources, there is a critical need to increase national capacity in coral reef science (including fisheries). Our advocacy on the impacts associated with climate change on the coral reefs and associated ecosystems can only be demonstrated through a systematic and consolidated efforts on collecting evidence based information on the changes to the various goods and services provided by these ecosystems. Coral reefs have resilience to natural environmental changes with their ability to recover and adapt over time. It is therefore important to identify such areas through systematic monitoring and extending the existing monitoring programmes including additional sites throughout the country.

M Shiham Adam, PhD
Director General

Acknowledgements

Captain Shareef and crew of the M. V. Mohara, Captain Anwar of M. V. Gaaviya and Captain Adam of M. V. Noah provided safe and comfortable logistical platform for our field trips. Captain and crew of M. V. Carina provided additional logistical support for N. J. Quinn during a September cruise. N. Thoufeeq, H. Amir, and Y. Rilwan assisted in the data collection and data entry. A. Chan and I. Abid assisted with Addu atoll surveys. B. K. Kojis and M. A. Abdulla assisted with the Maniyafushi reef survey training and reef survey. M. Shafiya and S. Naeem participated several of the training dives. Sham'aa A. Hameed, Shaffa A. Hameed and K. Kulak provided diving assistance on South Male' roving dives. Their assistance and good cheer helped expedite the expeditions.

R. Kikinger, University of Vienna and V. Wiesbauer-Ali, Water Solutions Pvt. Ltd., graciously contributed access to their excellent temperature time series data from Kuramathi reef, Rasdhoo atoll, Kandholhudhoo reef, Ari atoll and Huvafenfushi, North Male' atoll, respectively. G. Stevens and R. T. B. Moore kindly contributed coral reef survey data from Four Seasons Resort at Landaa Giraavaru.

N. J. Quinn is grateful to the Council for International Education for the J. W. Fulbright fellowship during the period of the study and for the hospitality and fellowship offered by Dr. M. S. Adam and his team at the Marine Research Center, Ministry of Fisheries and Agriculture.

A. A. Hakeem and his team of marine biologists at Banyan Tree and Ansanga Resorts and Spa kindly provided data on their coral eating starfish control program and hosted N. J. Quinn in visits to the resorts. S. Ali and G. van Weet are gratefully acknowledged for both their hospitality at Alila Villas Hadahaa and willingness to share their observations coral reefs.

We gratefully acknowledge B. L. Kojis for her observations and comments on the recovery of the reefs and for carefully reviewing an early draft of the manuscript.

Report Objectives

In this report we examine a commonly accepted suite of biological indicators of coral reef health from a geographically diverse set of lagoon, shallow Maldivian reefs from northern Haa Dhaalu to Addu atoll, south of the equator, based on field work undertaken from February – October 2009. The Marine Research Centre's (MRC) National Coral Reef Monitoring Survey (NCRMS) data set will provide a historical record, along with other outside agency studies, for assessing the recovery of Maldivian coral reefs to natural disasters in the past 12 years.

The annual sea water temperature variation for the Maldives will be discussed in the context of its relationship to geographical and seasonal variation and coral bleaching.

The status of the reefs will be broadly compared with the situation on other regional reefs that experienced bleaching.

Introduction

The Maldives is comprised of about 1192 islands with a 300 km² land mass and an exclusive economic zone of 859,000 km². Within atoll lagoon and reef areas comprise approximately 21,300 km². The 26 atolls (with 20 administrative atolls) lie along a north-south axis between 7° 06' 35" N to 0° 42' 24" S latitude and from 72° 33' 19" E and 73° 46' 13" E longitude in the central part of the Chagos – Maldives – Laccadive ridge in the central Indian Ocean. Subject to high amplitude sea level fluctuations during the Pliocene – Pleistocene with alternative periods of exposure and vertical reef growth the atolls presently rest on as much as 3 km of limestone overlying an Eocene volcanic basement (Aubert and Droxler, 1992). The islands formed 5500 - 4500 years ago (Kench, et al., 2005) during a period when sea level rose about 0.5 m above the reef tops.

About 193 of the islands are inhabited by native Maldivians, 93 islands have tourist resorts, and 55 islands are reserved for industrial and agricultural use. Only 33 of the inhabited islands have a land area of >1 km² and only three islands have an area >3 km². Coral reef statistics produced by the United Nation Environment Programme (UNEP) in 2003 ranked the Maldives as having the seventh largest coral reef system within its territorial boundaries. The Maldives is estimated to contain 3.14% of the total coral reef area of the world (United Nations Environmental Programme, 2003).

A wet monsoon dominates the period April to November when the winds blow from the southwest. A dry season occurs from December to March with lighter winds from the northeast (Kench, et al., 2006). The total population is about 299,000 people with 35% resident in the capital, Male'. About 80% of the islands have an elevation of <1 m above the mean high water mark (Ministry of Planning and National Development, 2008).

Clearly the health of the coral reefs, which are the physical basis for the Maldives, are of national strategic concern. President Nasheed called the plight of the Maldives to the world in his September 2009, address to the United Nations. He stated that failure to come to a meaningful agreement in the International Convention on Global Climate Change would be suicidal for the Maldives

The historically low population densities in the Maldives resulted in a large area of coral reefs where the influences of humans were low. For many centuries Maldivians preferred to eat pelagic fish instead of reef fish (Sheppard and Wells, 1998) so fishing impacts on reef fish populations were slight. Mining of the reef for rocks and sand for building materials depleted many lagoons and faros of corals (Brown and Dunne, 1988) and reduced fish biomass (Dawson Shepherd, et al., 1992). However, human impacts on reefs are increasing. Recently, nutrients from the capital city/ island Male' (Risk, et al., 1994), snorkeler and diver effects (Allison, 1996) combined with shark fishing, aquarium fish and sea cucumber collection (Adam, et al., 1997) and harbour construction have been changing the reefs.

Although the Maldives is completely founded on the remains of living coral reefs, it has only been towards the end of the 20th century that systematic quantitative surveys of reefs, from the northern atolls to the southern most atolls were conducted by the government of the Maldives through the Marine Research Center (MRC), Ministry of Fisheries and Agriculture.

Initially, surveys were limited to individual atolls, e.g. the notable descriptive work was mainly done around the British air base at Addu atoll (Stoddart, et al., 1966; Spencer-Davies, et al., 1971). On the “Xaifa” expedition at Fadhippolhu, Rasdhoo, Ari and Addu atolls, Georg Scheer conducted some of the earliest detailed “coral sociology” studies, using the plant sociology techniques of line intercept and quadrat methods. These techniques, introduced into coral reef research by Scheer (1971; 1972; 1974), became standard for benthic reef surveys. Scheer reported coral cover ranged from 50 – 80% from his observations from Addu, Ari and Fadhippolhu atolls. Along with the descriptive work of the “Xarifa” expedition (Hass, 1961; Wallace and Zahir, 2007), there were anecdotal reports of luxuriant, diverse reefs from divers dating back to the 1950s. The health and beauty of Maldivian reefs was further confirmed during the establishment of the multimillion dollar tourist industry when scores of dive schools were included in resort developments to cater to the diver seeking to “dive in one of the last paradises on earth.”

Pichon and Benzoni (2007) reported that there were 248 coral species from 57 genera, although they stated that the taxonomic status of some of the earlier described species were doubtful. However, Carpenter, et al., (2008) considers that according to the International Union for Conservation of Nature (IUCN) Red List Categories and Criteria: 1% -10% of the total species in the Maldives are Critically Endangered and endangered species, 10% - 20% are Critically Endangered, Endangered, and Vulnerable and 40% - 50% are threatened or near threatened. The threats are from both global climate change and local impacts.

Initial Reef Check Surveys and the 1998 Bleaching Event

Although there were previous surveys that MRC staff participated in, quantitative surveys of the coral reef community organized by MRC began in 1997 using the Reef Check coral reef monitoring protocol (Hodgson, 1999). These surveys yield a scientifically based estimate of several parameters commonly used to assess coral reef community health and resilience.

The unprecedented coral bleaching in 1998 around the world (Wilkinson, 1998) including the Maldives, resulted in extensive mortality to scleractinian, reef building corals. During this event the living coral animals expelled their commensal living plant called zooxanthellae exposing the unpigmented coral tissue and underlying skeleton. Prior to 1998 an isolated bleaching event occurred on the three central atolls in the Maldives was reported (Wood, 1987). While the 1998 bleaching may not have been the first, it was the most intense and sustained in recent history in the Indian Ocean (Goreau, et al., 2000).




Figure 1 Comparison of NCRMS 150 m point intercept transect percent live coral cover data at three sites in 1997 before and immediately after the bleaching in April – May 1998 bleaching event (after Zahir 2002a)

The NCRMS surveys clearly detail the death of live coral in Anbaraa reef, Vaavu atoll, Thuvaru reef, Meemu atoll and the high mortality of corals in Maduvvari reef, Meemu atoll (Figure 1). Prior to the bleaching event the reefs were considered luxurious at Anbaraa (Patch reef) and Maduvvari reef, with about 50 - 70% coral cover. Thuvaru reef was a moderately developed reef with less than a 30% coral cover prior to bleaching. Fast growing branching corals, such as *Acropora* spp., suffered greater mortality than slow-growing massive corals (Zahir, 2002a). Virtually all of the colonies of *Acropora* spp. in the study sites had died (Zahir, 2006). Similarly the fast growing pocilloporids were highly susceptible to bleaching. In contrast agariciids and poritids were more resilient, although they too experienced a decrease in cover (Zahir, 2).



Figure 2 A sea cucumber (*Holothurian atra*) lies among the rubble of the devastated reef flat (2 m) at Vabbinfaru reef, September 2009

Other workers found the percentage of living coral cover on several shallow reef flats declined markedly immediately after the bleaching event. Coral cover was reduced, from around 22.5 – 70% pre-bleaching to 0 – 10% post-bleaching (Allison, 1999; Clark, et al., 1999). The reef flat at Vabbinfaru, North Male' atoll exhibited very high mortality (Hakeem, per. com.). The luxurious reef at this site gradually became nothing but rubble / sand (Figure 2) which increased the beach erosion on the island which the resort tried to mitigate by pumping sand from the reef flat, further destabilizing natural efforts for recolonization.

Tsunami Devastation

Coral reefs are critical to the Maldives as they maintain the integrity of atoll islands by providing erosion barriers and are sources of sand and rock. They are also a major attraction for the tourist industry.

In December 2004, another natural disaster impacted Maldivian coral reefs. Originating in an earthquake off the east coast of Indonesia, a tsunami struck the Maldives on 26th December 2004. In February 2005, joint Maldivian and Australian research team surveyed 124 reefs sites in seven atolls, covering about 170 km of reef margin (AusAid, 2005) looking for tsunami damage. They observed localized damage to reefs near inhabited islands from debris (from buildings and other island infrastructure) and sand that had smashed and smothered corals when it had been swept into the ocean. Some of the damage was extensive enough to alter the reef framework. They concluded that there was minor direct damage to coral reefs by the tsunami, but only observed live coral cover in the range of about 4 – 12% (Figure 3). Although the reefs were not particularly luxuriant, they were recovering from the 1998 bleaching.

The researchers considered that the 1998 coral bleaching event caused more overall damage to the reef community than the tsunami (AusAID, 2005). They found that the tsunamis did not cause significant damage to Maldivian coral reefs, but it was likely that the tsunamis slowed recovery from earlier damage caused by bleaching. The damage from the tsunami was far less than the mortality resulting from the 1998 bleaching event and the on-going human damage to the reefs caused by coral rock and sand collection from reef flats and the dredge and fill operations associated with coastal development (AusAID, 2005).

The tsunamis did focus attention on the need for better management of direct human pressures and inappropriate coastal development. They note that their assessment of the condition of the reefs was hampered by the lack of comparable prior quantitative surveys that covered the length and width of the country at several depths (AusAID, 2005).




Figure 3 Percent live coral cover reported by surveys undertaken in January – February 2005, a couple months after the December 2004, tsunami (after AusAID, 2005)

Erosion

After the 1998 bleaching event, it was postulated that waves and bioerosion would operate synergistically to dismantle the reef framework and erode the islands (Figure 4). Boring and grazing organisms might remove CaCO_3 faster than primary frame builders could accrete it (Zahir, 2002b) and waves would break apart the weakened live rock. As well, it was thought that a tsunami would further wash away the islands. It was hypothesized that the structural integrity of the reef would be determined by the extent and rate of these biogenic and physical processes. Should these processes exceed the very limited accretion rates of the post bleaching reefs, the reefs' framework could begin to disintegrate further exposing the islands to the forces of oceanic swells as recorded in the Seychelles (Sheppard, et al., 2005).

Contrary to popular perceptions of the fragility and vulnerability of atoll islands, Kench, et al. (2006) showed that the uninhabited islands of the Maldives were robust landforms that experienced relatively minor physical impacts from the Sumatran tsunami. Morphological and sedimentary evidence suggests that although tsunamis do generate both erosional and depositional signatures, they do not promote gross instability of islands. Tsunamis are unlikely to be important mechanisms of atoll island destruction if the sand reservoir is conserved during the tsunami (Kench, et al., 2006). Where the beach sand reservoir was depleted and total beach volume reduced it is uncertain if the reefal sediment production will compensate for this reduction. Without healthy coral reefs, the replenishment of the biogenetic sand necessary for the natural maintenance of these dynamic islands would not be available.

While that finding is reassuring for the long term survival for the Maldives, many resorts were constructed with the intent to provide sea side villas. Without sufficient understanding the dynamics of sand islands and their sand budget under conditions with a stressed reef, the setback are now insufficient and many resorts are under threat from erosion (Figure 5).



Figure 4 Trees fall into the sea as waves on Vaavu atoll sweep across the reef eroding Bodumohora Island, March 2009



Figure 5 During high tide with strong westerly wind the sea washes with several meters of the villas, North Male' atoll, October 2009

In an experiment in 2001 - 2002, Zahir (2002b) investigated effects of bioerosion on coral blocks at two sites in North Male' atoll. He found that there was a significant difference between the sites in the intensity of bioerosion at shallow and deeper depths. However, it was unresolved whether bioerosion would damage a reef's structural integrity before it recovered. The answer to that question had to wait several more years when the reefs were resurveyed.

Natural Resilience - Recruitment

In spite of the calamities that had happened to Maldivian reefs the results from recruitment experiments were encouraging (Clark, 2000). Recruitment of juvenile corals to artificial structures 10 months after the bleaching event showed that 67% of visible (≥ 0.5 cm diameter) recruits were acroporids and pocilloporids and 33% were from massive families compared to 94% and 6%, respectively, in 1990–1994 (Edwards, et al., 2001). Although the recruitment rate had declined, the potential for restoration through natural recruitment looked promising.

Similar post-bleaching dominance of recruitment by branching corals was also seen on nearby natural reef (78% acroporids and pocilloporids; 22% massive corals) (Zahir, et al., 2002). Zahir et al. (2002) concluded that in spite of the mortality associated with the bleaching, sufficient numbers of coral colonies survived to produce "viable supplies" of planulae to recolonize the reefs. Later from the perspective of seven years after the bleaching, it appeared that the patterns and rates of coral recovery were not recruitment limited (Zahir, 2006).

Not everyone thought the reefs would recover. From their keyhole perspective at Komandoo house reef, Komandoo Island, Lhaviyani atoll, Schuhmacher et al. (2005) saw the slow and scattered formation of new reef structure, being outweighed by the collapse of dead protruding colonies of tabulate *Acropora* spp. Six years after the bleaching, the formerly three-dimensional structure of the reef flat and upper reef slope was a leveled field of rubble, only partly consolidated by encrusting corals. From their perspective they saw cascading deterioration of the status of the reef for the future (Schuhmacher, et al., 2005). Similarly, at Vabbinfaru, North Male' atoll, the once luxuriant reef flat had been reduced to rubble which has still not recovered (Hakeen, pers. comm.).

National Coral Reef Monitoring Survey Sites

As Zahir (2000) accurately observed, the expanse of reefs in the Maldives are enormous and the human and logistical resources available for monitoring them are small. Over the years MRC has done its best to assess the status of the reefs from the far north atoll of Haa Dhaalu to the southern most atoll across the equator, Addu atoll with its limited resources. The sampling design was strategic and targeted specific atolls (Figures 6a – 6f) for the following reasons:

- Haa Dhaalu (northern most atoll and a regional development target)
- Male' (central atoll with intensive tourism and other commercial activities)
- Ari (central atoll with intensive existing tourism development)
- Vaavu (south central atoll with a community-based integrated island resource management project underway).
- Addu-Gaafu Alifu (southern most atoll and a target for regional development with an international airport).

Within each atoll three reefs were selected. They were chosen because they were considered to be representative of the reefs in the atoll and/or because they had been previously surveyed. Owing to the limitations associated with sampling in deeper waters and logistical issues associated with supporting divers in remote reefs, the quantitative surveys before 2009 were mainly conducted at depths less than 3 m. Surge from oceanic swells and safety considerations prohibited site selection on outer reef crests. The specific site coordinates are listed in Table 1.



Figure 6a-f 6a) Map of Maldives with numbered and colored areas showing NCRMS sites. 6b) Map of most northern NCRMS sites in Haa Dhaalu atoll, 6c) Map of central NCRMS sites in North and South Male' atolls, 6d) Map of central NCRMS sites in Ari atoll, 6e) Map of central NCRMS sites in Vaavu atoll and 6f) Map of most southern NCRMS sites in Addu atoll. Orange indicates islands. Only site names are shown in each atoll for clarity, yellow circles indicate general location of the transects




Figure 6b Map of most northern NCRMS sites in Haa Dhaalu atoll




Figure 6c Map of central NCRMS sites in North and South Male' atolls




Figure 6d Map of central NCRMS sites in Ari atoll




Figure 6e Map of central NCRMS sites in Vaavu atoll




Figure 6f Map of most southern NCRMS sites in Addu atoll

Table 1 National Coral Reef Monitoring Survey sites with geographic coordinates and atoll and regional classification of the sites. Sites sampled in 2009 indicated with X. Thuvuru, Anbaraa, and Maduvvari were sampled from 1997-1999

Regions	Atolls	Sites	Geographic Coordinates	2009
Northern	Haa Dhaalu	Hon'daafushi	6° 46.33' N 73° 07.95' E	X
Northern	Haa Dhaalu	Finey	6° 45.06' N 73° 03.56' E	X
Northern	Haa Dhaalu	Hirimaradhoo	6° 43.86' N 73° 01.31' E	X
Central	North Malé	Bodu Ban'dos	4° 16.00' N 73° 29.12' E	X
Central	North Malé	Udhafushi	4° 18.85' N 73° 30.12' E	X
Central	South Malé	En'boodhoofinolhu	4° 07.12' N 73° 28.04' E	X
Central	Ari	Fesdhoo	4° 00.02' N 72° 00.32' E	X
Central	Ari	Maayaafushi	4° 04.94' N 72° 53.19' E	X
Central	Ari	Velidhoo	4° 11.53' N 72° 49.32' E	X
Central	Ari	Kan'dholhudhoo	4° 00.15' N 72° 52.88' E	X
Central	Vaavu	Anbaraa	3° 26.11' N 73° 25.33' E	X
Central	Vaavu	Vattaru	3° 13.49' N 73° 26.00' E	X
Central	Vaavu	Foththeyo	3° 26.90' N 73° 45.27' E	X
Southern	Gaafu Alifu	Koodoo	0° 43.91' N 73° 09.29' E	
Southern	Addu	Hithadhoo	0° 35.13' S 73° 06.14' E	X
Southern	Addu	Gan	0° 41.32' S 73° 09.72' E	X
Southern	Addu	Vilingili	0° 40.45' S 73° 11.48' E	X

Survey Methods

In 2009, the NCRMS adapted the Reef Check protocol (Hodgson and Liebler, 2002; Hodgson, et al., 2006) for surveying the reefs. Reef Check was designed to assess the health of coral reefs and is quite different from other monitoring protocols. It only varies slightly from the previous techniques used in the NCRMS to monitor benthic habitat. Reef Check does include a lot more parameters and is a more comprehensive technique that can be used to compare Maldivian reefs with other reefs around the world.

Since its inception, Reef Check has focused on the abundance of particular coral reef organisms that best reflect the condition of the ecosystem and that are easily recognizable to the general public. Selection of these "indicator" organisms was based on their economic and ecological value, their sensitivity to human

impacts and ease of identification. Sixteen global and eight regional indicator organisms serve as specific measures of human impacts on coral reefs. These indicators include a broad spectrum of fish, invertebrates and plants that indicate human activities such as fishing, collection or pollution.

Some Reef Check categories are individual species while others are families. For example, the Napoleon humphead wrasse (*Cheilinus undulatus*) is the most sought after fish in the live food fish trade, whereas the banded coral shrimp (*Stenopus hispidus*) is collected for the aquarium trade. Both species are very distinctive organisms and are considered by Reef Check organizers excellent indicators of human predation. The Napoleon humphead wrasse has been protected in Maldivian waters since 1995 (Notice No: FA-A1/29/95/39) (24 June 1995)) and is not a good indicator in Maldivian waters. It is hypothesized that on reefs where these organisms are heavily exploited, their numbers are expected to be low compared to their abundance on unexploited reefs. The analysis of fishes will be dealt with in a subsequent report.

Reef Check teams collect four types of data;

- 1) a description of each reef site based on over 30 measures of environmental and socio-economic conditions and ratings of human impact;
- 2) a measure of the percentage of the seabed covered by different substrate types, including live and dead coral, along four 20 m sections of a 100 m shallow reef transect;
- 3) invertebrate counts over four, 20 m x 5 m belts along the transect;
- 4) fish counts, up to 5 m above the same belt.

Monitoring of the indicators is done along two depth contours 3 m (2 - 6 m range) and 10 m (>6 – 12 m range) below sea level. Along each contour, four 20 m long line transects were surveyed using a measuring tape. The substrate under every 25 cm of the tape was recorded. Previously, three 50 m long transects were used from 1997 to 2005 and the intercept length to the nearest cm for each benthic category lying under the tape was recorded (Zahir, 2006). Transects followed the designated depth contour. Transect start and end points were separated by a 5 m space. Digital photographs were taken along the transects to document the species composition and abundance. The images will be analyzed using CPCe V3.4 software and discussed in detail in a later report. Additionally, roving divers randomly swam along the transect line to a depth up to 30 m recording unusual animals, diseased corals, and dying corals.

Results

Reef recovery: Coral cover

The mean and standard deviation of the coral cover from the two depths for each site was calculated. The mean coral cover ranged from 7.5% at Hirmaradhoo to 59.4% at Fesdhoo (Figure 7). There is a lot of variability in cover throughout the Maldives. Only four sites had a coral cover >50%: Fesdhoo, Kandholhudhoo (both Ari atoll) and Vilingili and Hithadhoo (both Addu atoll). The reefs in the narrow depth range sampled also are quite variable in their cover with the coefficient of variation ranging from 4.0% at Vilingili to over 100% at Hirmaradhoo and Maniyafushi.

In the following sections we will discuss the coral cover for each of the atolls in more detail.

Haa Dhaalu Atoll

Coral cover on the northern reefs of Haa Dhaalu atoll was nearly 0% after the 1998 bleaching event (Zahir, 2000; 2006). Subsequent reef recovery was slow with coral cover only achieving 3 - 5% in 2005. However, since 2005, coral cover has been rapid. In 2009, coral cover ranged from 14% - 29% (Figure 8). This was collectively the lowest coral cover for all the atolls surveyed. The northern atolls of the Maldives are more

exposed to cyclonic tropical storms than other atolls in the Maldivian chain and impacts by storms could setback recovery on northern reefs. No large storms impacted these atolls from 1998 to 2009.




Figure 7 Mean (3 m, 10 m) percent coral cover of the 17 NCRMS sites surveyed from March to October 2009. Error bars show standard deviation. n = 8 for each site

At 3 m at Hirimaradhoo, the live coral cover was only 14% - lower than at Hon'daafushi and Finey (Figure 9). Over 65% of Hirimaradhoo was covered with the category "other living organisms" which was mainly the invasive corallimorpharian, *Discosoma c.f. rhodostoma*. The remaining 20% was non living substrate. The corallimorpharian dominated much of Hirimaradhoo reef particularly at the depths of 3 -15 m. (Figure 10). It also was dominant in many Sri Lankan reefs post bleaching (Rajasuriya and Karunarathna, 2000).




Figure 8 Percentage live coral cover at 3 m at three sites in Haa Dhaalu atoll. Error bars only included for 2009, n = 4 for each site

Hon'daafushi and Finey had higher coral cover at 10 m than at 3 m. Hon'daafushi had a coral cover of 50% while Finey had 21% coral cover (Figure 9). At Hirimaradhoo because *Discosoma c.f. rhodostoma* was dominating the reef, hard coral cover was the least of the Haa Dhaalu sites with only 14% cover at 3 m and 0.6% at 10 m.

At 10 m hard coral cover at Hirimaradhoo was <1% because corallimorphirians dominate 80% of the reef cover (Figure 10), 1% algae and non living substrate cover 18%. The cover of corallimorphirians increased from 11% in 1998 to 40% in 2005. In 2009, their cover had increased to a 65%, overgrowing coralline algae, sponges and crowding out many other taxa on the reef, and diminishing space for coral settlement. This is the only site with a documented "phase shift" from a crustose coralline algae/ coral community to a corallimorpharian community. However, large beds of corallimorphs, covering >60% of the substrate, were recorded in 2003 at Addu atoll (Ministry of the Environment, Energy and Water, 2006).





Figure 9 Comparison of % coral cover at 3 m and 10 m at three sites at Haa Dhaalu atoll, March 2009. n = 4. Standard error shown by bars

Scleractinian corals compete with each other and other sessile organisms like macroalgae for settlement space and light using a number of competitive mechanisms including direct competition. The results of interactions between corals and macroalgae depend on the specific competitive abilities of different coral and algal species. In general, there is no clear competitive dominance of one group over the other (McCook, et al., 2001). Subsequent to large a die-off of corals, when unchecked, fleshy macroalgae have been known to increase in biomass and cover the substrate. There has not been a documented case where the reef changed from a scleractinian dominated substrate to a corallimorpharian dominated reef.

Figure 10 Percent cover of the corallimorpharian population *Discosoma c.f. rhodostoma* on Hirimaradhoo reef at 3 m from 1998 to 2009. Standard error bars only shown for 2009 (n = 4)



This change in benthic cover has been term “phase shift”. The term “phase shift” was first used to describe a change to Jamaican coral reefs after a series of devastating natural events (Hughes, 1994). After Hurricane Allen in 1980 and Hurricane Gilbert in 1988, the reefs off Discovery Bay, Jamaica exhibited a dramatic decrease in coral abundance and diversity (Woodley, 1989). Live coral cover dropped to almost 0%. Wapnick, et al. (2004) extracted cores from the reef and found that the transition from an acroporid dominated coral reef to a macro algal dominated reef was unique on a centennial or millennial scale. Wapnick, et al. (2004) considered that it was likely that the Jamaican reefs were exhibiting a community structure unique in the past several thousand years caused by extreme levels of both human and natural disturbance. Overfishing had devastated the benthic piscivorous and herbivorous fish populations. With the absence of *Diadema antillarum*, a sea urchin that had been devastated by disease, there were too few herbivores and algae covered the Jamaican reefs, drastically decreasing the amount of suitable substrate for coral planulae to settle and grow.

Healthy stocks of herbivorous fish were observed in Haa Dhaalu atoll (Figure 11). Maldivians have not intensively fished reef fish stocks, preferring to eat pelagic fish (Sheppard and Wells, 1988) leaving reef fish stocks relatively unexploited.



Figure 11 Schools of convict surgeon fish, *Acanthurus triostegus*, at 5 m, Hon'doofushi, Haa Dhaalu, March 2009

I is uncertain what controls *D. c.f. rhodostoma* population dynamics at Hirimaradhoo reef. Although they are soft bodied, we have never observed fish feeding on them nor do they exhibit feeding scars. Further research is needed to understand how Maldivian reefs respond when the hard corals die and new coral recruitment is insufficient to rapidly reestablish high coral cover. Have some reefs in the Maldives always exhibited high cover of corallimorpharians or is this truly a unique “phase shift” brought about by the opening of so much bare substrate for settlement of larvae of corallimorpharians after the 1998 bleaching event?

Subsequent to the previous survey in 2005 a small channel has been constructed to assist vessels across the Hirimaradhoo reef to the beach (Figure 12). While there is an obvious alteration of the reef framework by the construction, the affects appear to be isolated to the area adjacent to the channel. It is unlikely that this relatively small reef modification is causing such a profound change to a large area of the reef in the survey transect.



Figure 12 A boat channel was constructed through the Hirimaradhoo reef after the 2005 survey

North and South Male' Atoll *Maniyafushi*

Maniyafushi is the future research station of the Marine Research Center (MRC) and had not been previously included in the National Coral Reef Monitoring program. The island is classified as an industrial island and was previously used by a fishing company which left the island in 2008. The coral reef survey by MRC was conducted in February 2009. The survey site is along the reef in front of the island's jetty. Debris tossed from the jetty litters the sea bottom including batteries, construction debris and general trash (Figure 13). It is the most littered reef we surveyed.



Figure 13 Debris tossed from Maniyafushi jetty covers part of the reef, February 2009

The live coral cover is very high along the reef crest 63% (Figure 14). The shallow reef is dominated by acroporid and pocilliporid corals (Figure 15). Live coral cover diminishes to 24% at 12 m. At 24 m the substrate is mostly sand. The shallow reef has a much greater coral cover than at 10 m possibly because the transect line at 10 m was on a reef wall. Coral growth on vertical walls is inhibited by shading and when coral branches are broken or corals are dislodged, gravity often carries them into deeper water. However, in silty environments, corals on walls are seldom smothered by sediment.

No diseased corals were observed. However, it was common to see a recently killed coral during roving dives (Figure 16). It is difficult to attribute the cause of the death, but predation is probable either by a coral eating gastropod (Figure 17), or the pin cushion starfish, *Calcita schmideliana* (Figure 18). An occasional *C. schmideliana* was observed on the reef, but never actively feeding on coral. The coral eating gastropod, *Coralliophylia sp.* is recorded from the Maldives, but the coral eating gastropod is not *Drupella cornus*. Further effort is necessary to identify the gastropods that are eating the coral colonies.

Skeletons of the pipe organ coral, *Tubipora musica*, were scattered around the island in February 2009. The pipe organ coral is the only coral not banned from export. It is exported to India. Live colonies were not seen in any of the CRMS surveys.




Figure 14 Percent live coral cover at 3 m at five sites in North and South Male' atoll. Standard error bars only included for 2009. n = 4 for each site



Figure 15 M. A. Abdulla and B. L. Kojis swim past healthy *Acropora* colonies as they return from conducting a survey at Maniyafushi reef, February 2009



Figure 16 Recently killed juvenile *Acropora sp.* coral. Possibly killed by *Culcita schmideliana* or a coral eating gastropod




Figure 17 A swarm of coral eating snails have eaten an *Acropora sp.* colony



Figure 18 The pin cushion starfish, *Culcita schmideliana*, on reef. This starfish eats small corals

En'boodhoofinolhu

En'boodhoofinolhu had the lowest live coral cover in NCRMS in 2009. The reef has been used for sand mining for several decades. The sand is used in the construction industry in Male'. The mining is done from dhonis using divers who scoop the sand into plastic fiber bags which are then loaded on the dhoni and transported to Male'.

The sampling site is on the lagoon side of the reef and consequently receives the downwind sediment plume from sand mining operations. Live coral cover was low at 10.5% (Figure 14), but had risen from <3% during the period 1998 - 2002. Non living substrate (e.g. rock, rubble, recently killed coral and sand) accounted for >85% of the substrate. The remainder of the bottom was covered by an occasional sponge and fleshy algae.

Several *Acropora* colonies had broken branches and gaps where branches had previously existed (Figure 19). The branches at these damaged parts of the colony were either dying or dead, but the remainder of the colony appeared alive and healthy. The cause of this damage is unknown.

Bodu Ban'dos

The coral cover at Bodu Ban'dos after the 1998 bleaching was similar to many other reefs - a marginal 2% cover. In the years to 2002 the cover progressed at an uneven rate to 6.9%. Subsequent to the 2002 survey, construction of a harbour for the Bandos Resort occurred adjacent to the survey site. Regrettably, there was no survey immediately during the construction, or immediately after. Consequently, it is impossible to assess the effect the harbour construction had on the reef. However, despite the nearby construction the coral cover increased to 41% in September 2009. This is a remarkable recovery considering the proximity to a busy harbour with its constantly resuspended fine sediment. The recently killed coral is 1% - 2% in both the deep and shallow transects. This level is common for many of the reefs in the NCRMS in 2009. The coral community only has a few fleshy algae as the total percent of nutrient indicating algae for both transects is less than 1%.

Udhafushi

Udhafushi was the outer reef slope of a faru in the center of North Male' atoll. From the post bleaching coral cover of 1.3% in 1998, the site recovered slowly to 2.9% in 2002. While the percentage increase of 223% looks impressive the actual amount of additional hard coral cover is only 2.40 m over a 150 m transect. From 2002 to the survey in October 2009, the coral cover increase dramatically to 21% at 3 m and 18% at 10 m. However, the percent coral cover was still the least for North Male' atoll. Although the reef is in the center of the atoll and presumably far from oceanic swells, there was evidence of 5 m sections of the reef sliding down the reef slope.

Depth Variation

At Manyiyafushi and Bodu Ban'dos the coral cover was greater along the shallow transects than on the deeper ones (Figure 20). The reef surveyed at 10 m at Manyiyafushi was along an east west wall which receives less direct light during periods of the year. Consequently, one would expect less coral cover. It is likely that owing to increased turbidity associated with harbour sediments the substrate at 10 m at Bodu Ban'dos receives less light and therefore supports less coral cover. The coral cover at Udhafushi was only slightly lower at 10 m (Figure 20). Considering the variability of coral cover on the reef there was essentially no difference in cover between the two transects.



Figure 19 Damaged Acropora at En'boodhoofinolhu reef, South Male' atoll, at 10 m, September, 2009

At En'boodhoofinolhu reef the coral cover was greater at 10 m than at 3 m (Figure 20). Perhaps currents help to remove some of the silt before it smothers the deeper corals or the sediment settles in the shallower water and reduces coral grow and smothers early recruits. There are no studies of current patterns from the Maldives. This is certainly a potential area for future research to understand circulation patterns.




Figure 20 Comparison of % coral cover at 3 m and 10 m at one site at North Male' atoll and two sites at South Male' atoll, in 2009. n = 4 at each site and each depth. Standard error shown by bars

Ari Atoll

Fesdhoo, Maayafushi, Velidhoo reefs

The shallow reefs of Fesdhoo, Maayafushi, Velidhoo in the NCRMS at Ari atoll exhibited exceptional resilience. In 2005, most of these reefs had coral cover less than 11% (Figure 21). Fesdhoo reef exhibited resilience throughout the survey period. After the bleaching event in 1998, it quickly recovered to 10% coral cover in 2000 and then by 2005 had over 30% coral cover. In 2009, it was among the most luxuriant reefs surveyed with about 65% coral cover. Velidhoo reef also demonstrated remarkable recovery from about 11% in 2005 to 50% in 2009. Sometime after the sampling in 2005, the hotel at Fesdhoo reef had expanded constructing water bungalows over the reef flat, within 100 m of the reef edge. It does not appear that the sedimentation associated with the construction of the bungalows greatly thwarted the recruitment of new corals by either smothering new recruits or covering suitable substrate with silt rendering then unsuitable for settlement. Much of the recovery in these sites is due to successful recruitment and growth of primarily acroporid colonies and to lesser extent pocilliporid colonies. Maayafushi reef also had good recovery from 6% in 2005 to 31% coral cover in 2009.

Dega Giri

Surveyed in 2008 by the Marine Conservation Society, Dega giri in Ari atoll is among the reef with the highest coral cover (68%) (Solandt and Wood, 2008) (Figure 21). The substrate was dominated by mature colonies of *Acropora* with the coral cover at the giri edge being the greatest (Solandt and Wood, 2008).

Surveys using a standardized protocol, such as Reef Check, from various overseas organizations are important as they contribute to the knowledge of the conditions of the vast Maldivian reef system. It is however, suggested that MRC be designated as the official Reef Check coordinator for the Maldives and that all registered Reef Check groups coordinate their efforts through MRC and provides copies of their data set to MRC. On occasion surveys by foreign researchers are undertaken and little of the knowledge gained is retained in the Maldives.




Figure 21 Percentage live coral cover at five sites in Ari atoll. Standard error bars only included for 2009. (n = 4 for each site). Dega giri data from Solandt and Wood (2008)

Kandholhudhoo

Kandholhudhoo reef was added to the NCRMS in 2009. With over 50% coral cover it is among the more luxuriant coral reefs in the atoll. The island has been modified to be used as a picnic island by guests from nearby resorts (Figure 22). Presumably it was chosen for that purpose because of its nice sandy beaches and diverse coral reef (Figure 23) for snorkeling and diving.



Figure 22 Kandholhudhoo is a picnic island with beautiful beaches naturally replenished by a diverse, healthy coral reef which surrounds the island. April 2009



Figure 23 Kandholhudho reef at 7 m showing some physical damage and partial mortality to tabulate *Acropora* and intact tabulate *Acropora* colonies, March 2009

In Fesdhoo, Maayafushi, and Velidhoo the percent coral cover was greater in the shallow water than at 10 m (Figure 24). At 10 m, Kandholhudho reef had a 64% coral cover and was as luxuriant as the shallow Fesdhoo reef. Water bungalows have been built at Fesdhoo near the survey site since the last survey and have not adversely affected the reef.

The Velidhoo reefs here had the second lowest % coral cover at both the 3 m and 10 m surveys. It is unclear why the coral cover was so low. Maayafushi had the lowest coral cover (32.5% at 3 m and 13.8% at 10 m). The presence of a sewage outflow from the resort at the Maayafushi NCRMS site at 10 m is hypothesized as a causal agent in diminishing the ability of the reef to recover. This site would be an interesting research site to assess the effects of point source pollution on a remote and relatively pristine reef.




Figure 24 Comparison of % coral cover at 3 m and 10 m at four sites at Ari atoll, April 2009. n = 4 at each site and each depth. Standard error shown by bars

Vaavu Atoll

The NCRMS established a shallow water survey site at Anbaraa in 1997, before the 1999 bleaching event. This was among the most luxuriant reefs with near 60% coral cover (Figure 25). In a resurvey of the site shortly after bleaching in 1998, the coral cover had been dramatically reduced to about 1%. The acroporids and pocilloporids had died. Only a few massive (*Poritidae*, *Faviidae*) and non branching corals (*Agariciidae*) remained.

In April-May 1999, McClanahan (2000) surveyed 19 sites at a depth of 1 - 30 m (mostly in the Vaavu atoll region) and noted that the benthic cover was dominated by coralline and turf algae (68%), followed by erect algae (9%), hard coral (8%), sand (7%) and other invertebrate groups <3.5%. In 1999, *Porities*, *Astreopora*, *Pavona*, *Favites*, *Fungia*, *Favia*, *Leptastrea*, *Coscinarea*, *Symphyllia*, *Montipora* and *Goniastrea* accounted for 45.4% of the live coral in 1999, while *Acropora* accounted for only 3.5% of the coral cover. In March 2009, *Acropora* dominated the shallow water coral community at the three NCRMS sites. Two months after the December 2004 tsunami the reefs were resurveyed and compared to the 2004

NCRMS observations. In two of the three sites the coral cover had declined: 1) from 10.8% to 9% at Vattaru and 2) from 4.4% at Fohtheyobodufushi to 1.8% (Zahir, et al., 2006). The decline in coral cover was caused by sediment deposition and smothering of the corals by sand and debris that had washed across the reef flat (Zahir, et al., 2006). Both sites were lagoon side reefs on the eastern side of the Maldives facing the direction of the tsunami. No effects of the tsunami were observed at Anbaraa. This is possibly because the site is located in the middle of the atoll and because the reef is not orientated to waves sweeping over the reef flat.

In 2009, the reefs of Vaavu atoll have recovered from a percent coral cover of <10% in 1999 to a coral cover of 18% - 48% in March 2009. The reef by the most eastern island in the Maldives, Fohtheyobodufushi, had the highest coral cover for the atoll 48% (Figure 24). The island was uninhabited and was only occasionally used by fishermen and visitors from safari dive boats. On the other side of the atoll the reef at Anbaraa had increased from about 5% coral cover in 2005 to about 18%. To the south of Vaavu atoll is the small atoll of Vattaru. The reef surrounding Vattaru island is a Marine Protected Area, but is used illegally by fishermen to catch marine resources. The shallow reef coral cover had varied from 1998 to 2005 when coral cover ranged from about 2% in 1998 to 5% in 2003 before dropping to below 2% in 2005 (Figure 25). This reduced coral cover was due to sedimentation and subsequent smothering of the coral community due to 2004 tsunami as reflected by post tsunami assessment of reefs in February 2005 (AusAID 2005). A similar reduction in coral cover was recorded at Fohtheyo. These two sites were the only two sites of NCRMS that allowed to assess and compare pre and post tsunami impact to the coral community. However, by 2009, coral cover had increased dramatically to 32% and 48% at Vattaru and Fohtheyo respectively.




Figure 25 Percentage live coral cover at three sites in Vaavu Atoll. Standard error bars only included for 2009, n = 4 for each site and year. 1999 point represents pooled values from McClanahan (2000)

The percent coral cover was higher in shallow depths in two of the sites (Figure 26). Only at Anabarra was coral cover at 10 m greater than at 3 m. Additional abiotic data needs to be collected in order to further understand the spatial and depth variation within the atoll.




Figure 26 Comparison of percent coral cover at 3 m and 10 m at three sites at Vaavu atoll, May 2009. n = 4 at each site and depth. Standard error shown by bars

Gaafu Alifu Atoll

The single NCRMS site in Gaafu Alifu atoll, Koodoo, was only surveyed in 1998, 1999, 2002 and 2005. By 2005, there had been a dramatic increase in coral cover from about 6% in 2002 to nearly 26% (Figure 27). The increase is attributed to a rapid growth in tabulate acroporids (Zahir, 2006).




Figure 27 Percentage live coral cover at 5 m at Koodoo, and Han'dahaa in Gaafu Alifu atoll. n = 4 for Koodoo for each year. Han'dahaa based on visual estimate survey

In September 2009, Han'dahaa reef (0° 30' 19" N; 73° 27' 14" E) was inspected and coral cover estimated in a roving diver survey lasting 1 hour in depths up to 32 m. The Han'dahaa reef has the greatest coral cover of all the reefs in the NCRMS. Large tabulate corals (*Acropora cytherea*) up to 3 m in diameter dominated the reef from 3 – 12 m. Coral cover exceeded 100% along scores of meters of reef at the 10 m depth contour (Figure 28). Large tabulate corals commonly covered other corals resulting in coral cover exceeding 100%. Even at 20 m the coral cover was about 90% (Figure 29).

This luxuriant coral reef community exists in spite of a four year construction program to build the 50 room Alila Villas Hadahaa. Great care went into the construction of the resort to preserve the natural vegetation and minimize alterations to the fringing coral reef in order to maintain the natural ambience of the island and minimize the environmental footprint of the resort. The resort was the first resort in the Maldives to adhere to the standards for EC3 Green Globe certification for 'Building, Planning and Design Standard' Green Globe, an international benchmarking and certification program for the travel and tourism industry, is managed by EC3 Global, a subsidiary of the Australian based research body Sustainable Tourism Co-operative Research Centre (the world's largest source of tourism research). Its standards are based on the Agenda 21 principles for Sustainable Development endorsed by 182 Heads of State at the United Nations Rio Earth Summit in 1992. In keeping with this environmental achievement the management intends to implement a long term marine monitoring program. Future monitoring programs by the resort is likely to include subsurface sea water temperature, growth of selected coral species, monitoring coral cover, stock assessment of groupers, emperors, lobsters and other reef associated indicator species. The Maldivian owners are avid scuba divers that are passionate about their desire to insure that the best management practices are applied to the island's habitats.




Figure 28 Over lapping of tabulate corals (*Acropora cytherea*) at 10 m at Han'dahaa, Gaafu Alifu atoll has created one of the few sites known where coral cover exceeds 100%. September 2009



Figure 29 Acropora dominated the north east reef slope at 15 - 30 m, Han'dahaa, Gaafu Alifu atoll, September 2009

While large areas of the Han'dahaa reef slope are healthy with an extremely high percent coral cover, the western reef flat showed signs of large waves several years ago. Tabulate corals over a large area away from the resort's western loading jetty were detached and overturned. In spite of being dislodged and overturned many of the tabulate corals remained alive and some were growing in their natural horizontal shape (Figure 30).




Figure 30 Dislodged and overturned Acropora corals on reef flat, 4 m, Han'dahaa reef, September 2009. Note corals are still alive and new horizontal growth is occurring on left coral

During the resort construction large expanses of the western reef flat was covered by a filamentous algae (Ali, pers. comm.). By September 2009, most of the filamentous algae was gone and only a few coral appeared have died (Figure 31). The installation of in situ data loggers to record nutrient levels, salinity and water temperature and a terrestrial meteorological station would begin to provide the context for understanding changes to the habitat.



Figure 31 S. Ali inspects filamentous blue green algae on a tabulate coral and recently dead tabulate coral (white area) on the Han'dahaa reef flat near the desalination plant outfall (see upper left hand corner of photo). 4 m, September 2009

Addu Atoll

The NCRMS was conducted in May 2009, in Addu atoll (Figure 32). As with the other reefs in the Maldives, the coral reefs of Addu atoll were subjected to the 1998 bleaching event (Allison, 1999; Clark, et al., 1999), but had the greatest recovery of live coral cover in sites distant from anthropogenic influences. After the 1998 bleaching, the reefs at Addu atoll still had some living coral. At 5 m the Vilingili reef had about 4% coral cover in 1998 increasing to 13% in 2002. In 2009, the coral cover had increased to 55% (Figure 33) in spite of its proximity of the development of resort water bungalows and beach restoration efforts. Recently killed coral was observed (Figure 33) at Vilingili reef, but only covered 0.3% of the substrate in the two transect lines.



Figure 32 Y. Rilwan is recording substrate cover on a transect line at Vilingili reef, Addu atoll. 10 m, May 2009

During the Vilingili survey in May 2009, the sediment plume from the resort's beach restoration occasionally clouded the shallow parts of the site. The reef has survived the 1998 bleaching better than many other reefs as there were still large colonies of the slow growing *Porites lobata* coral (Figure 35).




Figure 33 Percentage live coral cover at 5 m at Gan, Vilingili and Hithadhoo reef, Addu atoll. Standard error bars only included for 2009, n = 4 for each site and year




Figure 34 Recently dead coral at Vilingili reef, Addu atoll, May 2009. Algae are beginning to grow on portions of the dead coral



Figure 35 Large colonies of *Porites lobata* were common at Vilingili reef, May 2009

Previous surveys of Addu atoll from 2002 - 2004 included transects at 10 m as well as shallow surveys at Gan, Vilingilli and Hithadhoo. Regrettably, there were no quantitative surveys below 5 m of these reefs immediately post bleaching, so it is difficult to assess both the damage and recovery. However, the reefs during this period had 40 – 63% coral cover (Figure 36). The Gan site is juxtaposed to a shipping jetty and in 2009 was littered with debris. Although the jetty has been there since the British used the island as a military air base, the live coral cover in 2002 was 42%. However, in the seven years from 2002 to 2009 the live coral cover dropped from 32% to 14%. Without more detailed information it made difficult to determine a cause for the decline.

The other two survey sites had very good coral recovery. Vilingili reef remained within the standard error of the last survey in 2004 at nearly 60% live coral cover. Hithadhoo reef also remained stable with a live coral cover of 52% staying within the standard error of the previous survey. The Hithadhoo was a diverse healthy reef (Figure 37) that was included as a Marine Protected Area a few years ago after an intensive study of the land and marine environment (Ministry of Environment, Energy and Water, 2006).




Figure 36 Percentage live coral cover at 10 m at three sites in Addu atoll, May 2009. n = 4 at each site in 2009 and n = 3 in 2002 – 2004



Figure 37 I. Abid, a member of the MRC coral reef monitoring team conducting a survey along four 20 m transects at Hithadhoo reef, Addu atoll in May 2009

The % coral cover at both depths at Vilingili reef and Hithadoo reef are high and not significantly different between depths (Figure 38) ($P>0.05$). The coral cover at Gan reef is much lower and there is a significant difference in coral cover with depth as previously discussed.




Figure 38 Comparison of % coral cover at 3 m and 10 m at three sites at Addu atoll, May 2009. n = 4 at each site and each depth. Standard error shown by bars

During a resource assessment for the establishment of the Maldives Protected Area System Project the reefs surveyed around Addu atoll surveyed clear signs of hard coral community recovery (Ministry of Environment, Energy and Water, 2006). The recovery was dominated by Acroporidae species and a single species of Faviidae – Echinopora. The size of corals that recovered varied within the lagoon with the larger and subsequent old coral colonies located on the reef edge and slopes of the northern ocean side reefs. The assessment found that the area with the greatest percent live coral cover and diversity of coral included the reef where the Hithadhoo NCRMS site is.

It was recommended that this area be managed under a regime that prevents anthropogenic activities from damaging and degrading the coral populations (Ministry of Environment, Energy and Water, 2006). The assessment considered some of the major anthropogenic threats to the habitat to be:

- Pollution from human waste, petrochemicals and rubbish,
- Increased sedimentation from terrestrial degradation,
- Resource extraction, e.g. removal of sand, rubble and coral rock,
- Resource exploitation, e.g. overfishing of holothurians,
- Habitat alterations – cutting the reef to create boating channels,
- Habitat destruction – anchor damage by bait fishing vessels.
-

Coral Framework Status

The coralline algae / sponge association dominates much of the coral reef framework on a healthy reef. Lasagna et al. (2008) observed that the abundance of bare rock, sand and rubble derived from dead coral breakage was the result of the loss of 3D structure in Maldivian coral reefs and that this substrate smoothing is known to represent an impediment to coral recruitment (Loch, et al., 2005). This loose material can limit coral recruitment (Loch, et al., 2002; Lasagna, et al., 2006), as coral larvae preferentially settle on encrusting calcareous algae (Heyward and Negri, 1999) and frequently avoid rubble as a settlement substratum (Sheppard, et al., 2002). In 1999, McClanahan (2000) observed that 11 of the 16 coral families known from the Maldives had recently recruited to the 19 reefs they surveyed. Almost the half of the recruits were Agariciidae, with Acroporidae and Pocilloporidae being rare. However, from 2001-2002 the taxonomic composition of recruits shifted from a dominance of Agariciidae in the early stages of recolonization toward a dominance of Acroporidae and Pocilloporidae (Bianchi, et al., 2006; Zahir, 2006).

Roving diver surveys were conducted on the deeper reefs throughout the NCRMS in 2009. On many of these reefs (>20 m) Tubastrea micranthata was the most abundant coral. Acroporid and pocilloporid colonies were dominant in shallow waters (<10 m). Massive corals such as Porities, Pavona and faviids were usually small (<30 cm). These are slower growing taxa than acroporans and pocilloporids.

Colonies of *Porites* were observed with new growth on top of dead colonies (Figure 39). It is unclear whether this was re-growth of a surviving part of the original colony or new recruitment. This is certainly a question waiting for a future researcher to answer. If these colonies are survivors of the bleaching event, then it is likely that the colonies will survive future thermal increases. The comparative size of the *Porities* colonies against the other species (Figure 39) suggests that either they are survivors or early recruits to this open space.



Figure 39 Dead *Porites* colony with new colonies of *Porities*, *Acropora* and other species growing on it, 20 m, Magiri reef, North Male' atoll, September 2009

It is unclear whether there has been any local extinction of coral species. Bianchi, et al. (2006) reports that up to 2003, no colonies of *Stylophora* or *Seriatopora*, two genera previously reported in the Maldives prior to 1998, were observed at any site. Indeed, in this survey colonies of these genera were very rare. However, H.Z. has observed both genera on the Ari atoll outer reef slopes and suggests they are slowly returning in abundance.

A couple dead colonies of *Seriatopora* were for sale at Gloria Maries, a souvenir shop, in Male' in October 2009, for Rf4000 (Rf12.75 = US\$1.00). Maldivian law prohibits the export of scleractinian coral.

Only a few living remnants of *Acropora palifera* were observed post bleaching by Loch, et al., (2002) and by N.J.Q on the NCRMS. H.Z. has observed that *A. palifera* is common on western Ari atoll outer reefs. As well, *Millepora*, a common genus before the bleaching, was not found in surveys to 2003 (Bianchi, et al. (2006). Nor was it observed in 2009 in the NCRMS. In contrast, *Millepora* rapidly recruits in the Caribbean after bleaching (Kojis and Quinn, 2001).

Portions of colonies of *Porites c.f. lobata* and *Diploastrea heliopora* survived and grew 40 mm and 12 mm, respectively, post bleaching to March 2001 (Schuhmacher, et al., 2002). Perhaps the surviving remnants of these massive corals have developed a relationship with heat adapted zooxanthellae and are better adapted to increasing water temperatures. Following these colonies would yield interesting scientific information about a coral communities' adaptation to climate change.

While some habitats have recovered there are still large areas of reefs with little sign of recovery. Either the substrate is unsuitable for settling and survival or planulae are not being swept past the reef. The Guraidhoo Corner reef crest (8 m), South Male' atoll (Figure 40) and Vabbinfaru reef, North Male' atoll (Figure 2) are examples of two different habitats with poor recovery. Guraidhoo Corner is at the edge of a channel with strong currents and an abundant fish community. The area is a popular dives site because of the abundance of herbivores, groupers, sharks, turtles and manta rays. Vabbinfaru reef surrounds Banyan Tree Resort which pumps sand from the reef to replenish its beach which is periodically swept away by strong waves. An informed and concerned resort management understands the importance of the reef for the physical stability of the island and for its attraction for visitors. An innovative coral restoration program is endeavoring to assist in the recovery of live coral on the reef.



Figure 40 Guraidhoo Corner, South Male' atoll is a reef subject to oceanic swells and unconsolidated corals with poor recovery. September 2009, 8 m

While a sandy, shifting substrate may inhibit coral recruitment, for many decades Maldivians mined coral for building materials and the overall reefs seem to have survived. Historically, virtually all village homes, businesses and municipal building were constructed from coral. There were some shallow reef flats that were intensively and repeatedly mined resulting in these lagoons and faros being depleted of corals (Brown and Dunne, 1988). Legislation in the late 1990s recognized the environmental and economic damage of this practice and limited both the total volume and spatial extent of the coral removed. Coral mining is now regulated and specifically, coral cannot be mined on inhabited islands, resort house reefs, reef frequented by tourists, atoll rim reefs and common bait fish fishing reefs (Sluka and Miller, 1998). Many village homes now show a mixture of traditional coral and newer sand cement block construction.

The loss of complex physical structure can result in local extinctions, substantial reduction in species richness and reduced taxonomic distinctness associated with diverse productive coral reefs (Graham, et al., 2006). Many Maldivian atoll outer ocean facing reefs and the inner lagoon reefs (faros, thila or giris) do not have deep spur and groove features that characterize the Florida reef track and some southern Maldivian outer reefs. Many lagoon reefs exhibit steep slopes without major contours. In spite of the deterioration of the structural complexity of some reefs (Loch, et al., 2002) a large number of reefs have recovered very rapidly. At some sites there was a "tremendous increase" in hard coral cover shortly after the bleaching event (Loch, et al., 2002).

At a depth of 15 - 30 m caves are occasionally observed and are likely the result of erosion caused by wave action when sea level was lower during the last ice age which ended 8100 - 6500 years ago when sea level rose ~ 7 mm annually (Kench, et al., 2009b). These caves, formed by sea level still stands, have low light levels and consequently usually no zooxanthellate scleractinian corals. They attract low light level species such as Tubastrea corals, sponges, sea whips, black (Antipitherian) corals along with soldier fish and occasionally glass fish. There is no record of any surveys of these communities being conducted during or after the bleaching event. These are special communities that deserve particular consideration and monitoring.

Coral Reef Recovery in the Region

Dubai, United Arab Emirates

How have other reefs in the region recovered from the bleaching event? A decade after coral bleaching in the United Arab Emirates (UAE), one area formerly dominated by Acropora was dominated by faviids and poritids, with adult and juvenile composition suggesting this dominance shift is likely to persist (Burt, et al. 2008). Acropora dominated assemblages in the UAE were observed in three of the six sites examined and coral cover ($41.9 \pm 2.5\%$) was double the coral cover immediately after the bleaching. Porites lutea and Porites harrisoni dominated communities that were negligibly impacted by the bleaching events, and

the limited change in coral cover and composition in intervening years was probably from slow growth and low recruitment. Despite strong recovery of several dominant *Acropora* species, five formerly common species from this area were not observed suggesting local extinction. It was concluded that Dubai coral communities exhibit both resistance and resilience to elevated sea temperatures (Burt, et al., 2008).

Lakshadweep Islands, India

Closer to the Maldives in the Lakshadweep Islands, India, by 2005 the coral recovery was very site specific and was influenced by differences in post settlement survival of recruits which were driven by local water circulation patterns (Arthur, et al., 2006). In 2001 the coral cover ranged from 6% - 19% as the rate of recovery observed was not uniform. Recovery was greatest on the west facing reefs with very limited recovery on the east facing reefs. The reef did not experience a 'phase shift' to macroalgal dominated reefs. The presence of intense herbivore grazing was considered to be the most important controlling factor (Arthur, et al., 2006). The genera that showed the maximum increases five years after the bleaching were those with a mix of "different susceptibilities" to bleaching. Other genera that were not particularly vulnerable to the bleaching showed significant declines. The authors suggest that individual life history strategies, post-recruitment and circumstances greatly influence the decline and recovery of the coral communities (Arthur, et al., 2006).

Chagos Archipelago, Indian Ocean

To the south of the Maldives the Chagos Archipelago was particularly badly affected, suffering total or very heavy coral mortality on seaward slopes to >30 m depth with species-specific mortality extending deeper still (Sheppard, et al., 2002). This occurred in spite of its remoteness from direct human impacts. Hard and soft coral cover on seaward slopes before 1998 was 50% to 95%, which declined in 1998 to an average of 12%, and even to zero% at the depth of 0 – 5 m in some shallow areas (Sheppard, 1999). Initial mortality of corals was followed by a collapse of the reefs' structural framework resulting in the loss of obligate corallivores (Spalding and Jarvis, 2002).

As in the Maldives, fleshy algal cover did not increase after the coral mortality. By 2006, coral cover was almost restored to pre-1998 levels at most shallow sites, but had recovered much less in deeper waters. Sheppard, et al. (2008) considers that the coral cover alone is a poor indicator of recovery because the cover was dominated by *Acropora palifera* and other corals that are largely encrusting in juvenile form, in contrast to their mature morphology, in which they have a great 3D relief structure. Spatial complexity is important to diversity and abundance within the reef community, including fishes (Graham, et al., 2006) and needs to be considered when assessing reef recovery.

Coral Reef Diseases

No diseased corals were recorded in the NCRMS transects. However, a disease associated with crustose coralline algae was observed, but not quantified, during roving dives (Figure 41).



Figure 41 Coralline algae being attacked by the bacterial disease, CLOD (yellow), at En'boodhoofinolhu, North Male' atoll, September 2009. White section is recently killed coralline algae

Crustose coralline algae (Rhodophyta, Corallinales) are plants that deposit calcite, a particularly hard and environmentally resistant form of calcium carbonate. These marine plants bind surficial carbonate materials, debris, and other calcareous organisms to create stable substrata. Tropical reefs in high energy environment depend on calcareous coralline algae for the maintenance of wave resistant intertidal ridges (Littler and Littler, 1997). Many crustose corallines have a prostrate type growth form and are conspicuous as maroon, red, pink and purple pavement covering large areas, whereas other forms develop vertically branching heads much like some corals.

Functionally, crustose coralline algae, particularly *Porolithon*, are the principal binding agents that generate the structural integrity and shock resistance of the reef rim in high energy systems. Coralline algae are important for the absorption of massive wave energy that would otherwise erode shoreward land masses and for the facilitation of the development of many other sheltered shallow reef communities. All coralline algae proved fatally susceptible to the bacterial pathogen, CLOD (Littler and Littler, 1997). The bacterial disease should be included in future monitoring efforts.

Fish Distribution and Abundance

Using the Reef Check methodology at NCRM sites figures 42 a-e summarise the average fish abundance from the sites within each atoll. Figure 43 summarises the mean abundance of the various Reef Check fish categories. On average, Ari atoll had the most fish, followed by Addu, and Male' atoll had the lowest abundance. Butterflyfish consist of many species in Maldives and this was the main category driving variations in abundance. Many butterfly fish are dependent on or strongly associated with coral and so abundances and species richness can often be directly correlated with percentage coral cover. The results presented here follow this trend somewhat, with the two atolls having highest coral abundance, also having the most fish.

The sites in Ari generally showed a high abundance of fish. At Velidhoo, however, both at 5m and 10m, fish abundance was significantly less than at other sites. During reconstruction of the resort in 2005, water bungalows were added and it is possible that these may have had an effect on the resident fish populations. Coral cover, however was not greatly affected.

In Addu, Gan showed a drop in coral cover between 2005 and 2009, and had the lowest percentage coral cover of the sites in the region. This was also the site with the lowest fish abundance. It had significantly less butterflyfishes than at the other sites, however overall abundance remained high due to the presence of more parrotfishes. Villingili, where coral recovery and cover were highest had a significantly higher abundance of fish than the other two sites in the region.




Figure 42 a-e Average abundance of each Reefcheck fish category at the NCRM survey sites in each atoll

Haa Dhaalu had the lowest coral cover, however fish abundance remained relatively high. However, this high abundance was driven by large schools of five lined snapper (*Lutjanus quinquelineatus*). Butterflyfish and parrotfish were not seen in huge numbers. Interestingly, in Vaavu atoll, Vattaru is a marine protected area, and yet had the lowest fish abundance in the region.

Of all the regions surveyed in the NCRM programme, Male' atoll had the lowest fish abundance. A reason for this could be that this region has a far higher population than anywhere else in Maldives. Many reef fish, especially butterflyfishes are targeted for the aquarium trade and the proximity to the capital may make this the primary area for harvesting fish. Other reef fish have increasingly become subject to exploitation due to high demand from the resorts. Malé atoll also has a high concentration of resorts and fishing pressure nearby may be higher in order to meet the demand from tourists. Many resorts also offer reef fishing trips and these are rarely regulated.




Figure 43 Average abundance per category of fish in each atoll surveyed

Grouper Abundance

The grouper fishery in Maldives started in 1994 and has since become a valuable export. Slow growing and highly vulnerable to overexploitation, the Marine Research Centre has focused efforts to better understanding the status of the fishery and populations in order to improve management of the resource (see Sattar and Adam 2005). During the NCRM surveys, only 2 groupers over 50cm were encountered. These were in Vaavu and Haa Dhaalu atolls, although the highest abundance was in Ari atoll. The limitation of the Reef Check methodology in analysing the distribution of groupers, is that individuals under 30cm are ignored. For an exploited fishery such as this, it is essential to record even the small individuals. On some transects many small individuals were observed but not recorded. This suggests that groupers are recruiting successfully, and provides further evidence that the fishery is fully or over-exploited and that efforts researching the aquaculture potential and the fishery must continue if it is to remain productive.

Other Reef Fish

Overall, a low abundance of predatory reef fishes were encountered across Maldives. Grazers, on the other hand tend to be numerous and large in size. Compared to other countries, rich in coral reefs, Maldives does not use fish traps, and predators tend to be targeted mainly through bait or lure line fishing. This reduction in predators may be a reason for thriving grazer populations. The relatively low densities of *Diadema* urchins and low macro-algae cover support the notion that grazing pressure due to fish remains high on Maldivian coral reefs (NC pers. obs.). This was the first year that all sites included fish transects at two different depths and this should become a regular feature on the NCRM programme. Reef Check, although useful to give a summary of the status of fish populations, is not detailed enough to be able to monitor the combined influences of the reef fish fishery and the aquarium fish trade. It is suggested that more detailed fish surveys be incorporated into the programme to enhance current efforts in monitoring this fishery. For more information on the current status and recent management initiatives (see Sattar 2008). With the recent Darwin Initiative project recently launched at MRC, it is likely that in following years, the fish component of the surveys will become more detailed, with the information yielded giving valuable insight into the overall status of the coral reef ecosystem.

Sponge Abundance and Distribution

Overall non boring sponges represent <1% of the total cover in the 17 sites in the NCRMS. Overall the cover was: 3 m - 0.7% cover, 10 m - 0.7%. Non boring sponges were so uncommon that they were not present in transect line at Hirimaradhoo, Fesdhoo, Velidhoo, Kandholhudho, Udhaafushi, Maniyafushi, Gan, Vilingili and Hithadhoo (Figure 44). The greatest cover of sponges occurred at Anbaraa (6.3%) and En'boodhoofinolhu (5.6%).




Figure 44 Geographic and depth variation of % sponge cover in NCRMS sites in 2009

Terpios sp. (Demospongia, Order Hadtomrida) is an encrusting sponge which kills corals (Placer-Rosario, 1987) by getting nutrients from coral tissue. It is only about 1 mm thick but in the Maldives it has been observed by the roving diver surveys to cover many species of coral as well as competing for space with corals. The occurrence of *Terpios sp.* in the Maldives has not been previously reported in the literature. Although not specifically recorded in the sampling protocol, the sponge was seen overgrowing corals at Kandholhudhoo (Figure 45), one of the most luxuriant shallow water reefs surveyed. Kandholhudhoo is a picnic island used by resort guests. Kikinger (pers. comm.) reported the occurrence of *Terpios* in Ari atoll in 2005. However, in 2009, the occurrence of *Terpios* was rare at NCRMS, but future monitoring efforts need to identify it and minimally record its presence.



Figure 45 *Terpios sp.* overgrowing *Acropora* coral, Kandholhudhoo, Ari atoll, 10 m, April 2009

Echinoderms

Crown of Thorns starfish - *Acanthaster planci*

The Crown of Thorns (CoT), *Acanthaster planci*, and *Culcita schmidelina* starfish feed on live coral. In the late 1960s to the 1990s large swarms of CoT were observed feeding on the Great Barrier Reef and other Pacific reefs. Large areas of live coral on the GBR were killed and the CoT was seen as a major threat to the GBR's sustainability.

No CoT were observed at any of the 17 survey sites or during about 30 hours of roving diver surveys in the NCRMS in 2009. The only CoT observed was a juvenile CoT collected at Bodumohoraa Island, Vaavu atoll in 2008 (Figure 46).



Figure 46 Dead juvenile Crown-of-Thorns starfish, *Acanthaster planci*, from Bodumohora reef, Vaavu Atoll, October 2008

The presence of *C. schmideliana* was not part of the survey protocol and was not specifically recorded in the transects. However, typically at least one individual starfish was observed at each site. The occurrence of the starfish should be included in the protocol for future surveys.

The Maldives Protected Area System Project observed two adult COT at Addu atoll during their resource evaluation in 2003 (Ministry of Environment, Energy and Water, 2006). The report concluded at that time, the standing stock of COT in Addu atoll was no threat to the reefs.

A team lead by A. A. Hakeem collected both CoT and *Calcita schmideliana* several times a month from Vabbinfaru (4° 18' 35" N; 73° 25' 26"E) and Ihuru (4° 18' 24" N; 73° 24' 58"E) reefs in 2006 and 2007. In two years, a total of 84 CoT and 104 *C. schmideliana* were collected at Vabbinfaru ranging from 7 to 14 starfish per month (Figure 47). At Ihuru reef, a total of 66 COT and 84 *C. schmideliana* were collected ranging from 3 to 11 starfish per month (Figure 48). About 20% more starfish were collected at Vabbinfaru than at Ihuru reef in that period. In 2006, the team collected 44 CoT and 61 *C. schmideliana* from Vabbinfaru reef. The number collected in 2007 was 40 CoT and 43 *C. schmideliana*. At Ihuru reef in 2006, 37 CoT were collected and 47 *C. schmideliana* were collected. In 2007, 29 CoT and 40 *C. schmideliana* were collected.

Vabbinfaru reef is bigger than Ihuru reef. If each collection was from the entire reef one would expect more starfish to be collected at Vabbinfaru than at Ihuru reef. Also, since, there is insufficient information to standardize for collecting effort per month, we can only make the generalized observation that there appears to be a low level, persistent population of *A. planci* and *C. schmideliana* on Vabbinfaru and Ihuru reefs. It is also likely that the vigilant collection activities have helped to keep the population in check, although feeding scars are commonly seen on the reef.




Figure 47 Monthly records of *Acanthaster planci* and *Calcita schmideliana* collected from Vabbinfaru reef in 2006 and 2007 (Azeez, pers. comm.)

Echinoids

A total of 35 individuals of *Diadema antillarum*, 45 pencil urchins (*Heterocentrotus mammillatus*) and no collector urchins (*Tripneustes gratilla*) were observed from 17 sites and 13,600 m² of reef. The number does not increase when roving diver observations are considered as 98% of the urchins were observed in the shallow transect. *H. mammillatus* urchins were only recorded at the sites in Haa Dhaalu (Figure 47). *D. antillarum* was most abundant at Udhaafushi. It was not observed in 12 sites, included all sites in Ari atoll (Figure 49).




Figure 48 Monthly records of *Acanthaster planci* and *Culcita schmideliana* collected from Ihuru reef, North Male' atoll in 2006 and 2007 (Azeez, pers. comm.)




Figure 49 Abundance and distribution of two urchin species in the NCRMS sites in 2009


The Indian Ocean is not the Caribbean Sea and urchins play a different role in the reef community. Urchins in the Caribbean are important to clear space for coral settlement in a sea with only sparse herbivore fish populations. The absence of many urchins in the Indian Ocean does not correlate with the condition of the reef. Herbivores fish are not a target species and their large population function to keep fleshy algal levels low and space available for coral settlement.

Holothurians Abundance and Distribution

Relatively low densities of sea cucumbers were encountered at all NCRMS sites compared to reports and observations from other regions in the Indian Ocean such as Seychelles (NC pers. observ.) In Chagos, unexploited populations were at densities of 55.4 individuals per 100m² (Price et al. 2009; densities standardised for purpose of comparison in this report). The NCRM surveys revealed no significant difference in abundance at the different depths. Density varied between 0-3 individuals/400 Muthiga (2008) also reported a similarly low density along the reef edge of with 1.64 ± 1.04 individuals per 100m² in North Male' atoll. Figure 50 shows the overall abundance of sea cucumbers for each atoll. This includes the individuals seen during the NCRM surveys and also during free rover dives.

In all the regions sampled, *Personathuria graeffii* was the most abundant sea cucumber. Commercially, due to the small size of the body wall, this species is of negligible significance, although their importance to coral reef ecosystem health has been increasingly implied. The recently described *Bohadscia atra* (Mas-sin et al. 1999), which is still commonly believed to be *B. argus* was the most abundant commercial sea cucumber at the NCRM surveyed sites, with a maximum density of 2 individuals per 100m² at Velidhoo, Ari Atoll. Although harvestable, this species is of low-medium value. Ari atoll also had the highest species diversity with all of the species in table 2 recorded.

Figure 50 Total abundance of sea cucumber individuals spotted at each of the surveyed atolls including Reefcheck and free roving dives



The majority of medium-high value species were recorded in free rover dives at a depth >20m. *S. Hermannii* and *Thelenota anax* were found in all five regions, with the highest number of individuals of both species found in Ari Atoll. In the same location the highly valued and globally rare sea cucumber *H. nobilis* was seen. This was the only one encountered throughout all the surveys. Three individuals of the highly valued *Thelenota ananas* were seen in Haa Dhaalu (Figure 51).

Table 2 List of species encountered on NCRM surveys and during free rover dives at >20m with the commercial value for each species (adapted from FAO statistics, Conand 1990). X INDICATES species present in survey

Species	NCRMS	Free Rover Dives	Commercial Value
<i>Actinopyga miliaris or obesa</i> *	x	x	Medium
<i>Bohadscia atra</i>	x	x	Medium/Low
<i>Holothuria atra</i>	x		Low
<i>H. edulis</i>	x	x	Low
<i>H. nobilis</i>		x	Medium
<i>Pearsonothuria graeffei</i>	x	x	Low/ None
<i>Stichopus chlorontus</i>	x		Low/ None
<i>S. hermanni</i>	x	x	High
<i>S. pseudohorrens</i>		x	High
<i>Synapta maculata</i>		x	Low
<i>Thelenota ananas</i>		x	Medium/High
<i>T. anax</i>		x	High

*it is impossible to indentify between these species *in situ* without spicule analysis.

As this was the first time that sea cucumbers were included in the NCRM surveys, there is little historical data with which to compare distribution. Previous studies have focussed mainly on the fisheries management aspect (Gillet 2004, Ahmed et al. 1997, Joseph et al. 1992) or on the ecology of an individual species (Reichenbach, 1999). Additionally there is no data available prior to 1985, when exploitation of the resource commenced. Therefore it is unclear as to whether abundance and diversity are naturally low in

Maldives or whether exploitation has had a long term effect upon the natural populations. From the three studies cataloguing species encountered in Maldives (Muthiga 2008, Reichenbach 1999, James 1992) a total of 23 species have been recorded. The total number of species found in waters surrounding India is 79 (James 1992), and it is probable that there are more to be recorded in Maldives. A cumulative curve for species encountered in Muthiga's (2008) study shows a positive trend as dive time increases, which does not level off by the total dive time. It is recommended that a large scale study into the fishery and population status is conducted in order to maximise the outputs of this fishery, the demand for which continues to grow. As sea cucumbers are considered to be integral to coral reef health, acting as bio-turbators thus remineralising any settled nutrients into a useable form, it is doubly important that distribution and abundance patterns in Maldives are better understood.

Although historically there was no traditional sea cucumber fishery in Maldives, a huge surge in demand over the last three decades from China has meant that for many countries, Maldives included this has become a valuable fishery bringing an additional source of income to many people. Unfortunately the rapid pace of expansion of the fishery has meant that often the fisheries are unsustainably exploited and thus susceptible to collapse.

In Maldives, exports commenced with a trial batch of processed product of 31kg in 1985. By 1988, 553t were exported to a value of US\$4,496,327. By 1990, the fishery peaked for the first time volume wise with an export of 745t (MOFA statistics). Harvesting used to be done by hand collection, with a natural limit to the depth of collection. However, by the 1990s SCUBA was the most common method for harvesting, and deeper dwelling, valuable species were collected. Export figures indicate that by the early 1990s overexploitation of the resource was evident with less valuable species making up a greater proportion of the catch, and the price per kilo decreased. In 1993, MOFA banned the use of SCUBA for sea cucumber harvesting. Due to the large resources required, this ban has not been enforced and anecdotal evidence suggests that SCUBA is still the main method used for harvesting.

Further to this, because of the continued growth in demand for sea cucumbers, coupled with a susceptibility of these fisheries to collapse, technologies for culturing sea cucumbers continues to advance (for a review see Lovatelli et al. 2004). With the need for the economy in Maldives to diversify and decentralise away from Male', sea cucumber aquaculture is a possibility that should be further explored in the future.




Figure 51 *Thelenota ananas* at 20 m at Coral Gardens, North Male' atoll, October 2009

Crustacean Abundance and Distribution

Lobsters

The following four species of spiny lobster have been reported in the Maldives: *Panulirus penicillatus* (the doubled spine lobster), *P. longipes fermoristriga* (the long legged spiny lobster), *P. versicolor* (the painted lobster) and *P. ornatus* (ornate lobster) (Ahmed, et al., 1997). The lobsters occupy a wide range of habitats, however each species responds differently to habitat gradients, such as depth, turbidity, coral cover, wave action and feeding preferences (Wright, 1992).

P. versicolor is the most common lobster within lagoons and is found underneath table corals and in crevices. While it can be found in exposed reef slopes it is rarely found deeper than 6 m (Ahmed, et al., 1997). The species is also nocturnal and shelters in crevices during the day and forages at night. The species can be seen during the day as they are known to aggregate together with their long white antennae protruding from the crevice. Lobsters were first recorded in the NCRMS protocol in 2009. The previous field survey was Jonklass (1961).

In a total of 17 NCRMS sites covering five different atolls and a total area of 13,600 m² only four adult and one post puerulus lobsters were recorded. The number only slightly increases by eight lobsters when about 30 hours of roving diver observations are included. The low lobster population may be a consequence of few refuges associated with relatively low habitat relief and intense fishing pressure to supply resorts.

At Gan, Addu atoll, a post puerulus lobster (C.L. ~ 1 cm) was observed at 5 m among the rubble / coralline algae (Figure 52). No information is available about spawning seasons in the Maldives. This would be an excellent future research project done in collaboration with fishermen and resorts.

Lobsters are mainly caught by hand while free diving. Lobsters are not prized as a food for the local citizens. However, there is a constant demand by tourist resorts for fresh lobsters, so fishing pressure is high. The catch is either sold to local resorts or sent to Male' for re-sale. The catch statistics from MoFAMR for the period 1988 to 2005 ranged from 20,000 lobsters in 2005 to almost 70,000 lobsters in 2002 (Anderson, 2006). The resorts could assist in coral reef monitoring by recording and reporting number, weight and value of lobsters purchased. However, this would presumably only provide an estimate for adult male lobster abundance and distribution. The capture of female lobsters and those less than 25 cm in total length have been banned since 1993 (No: FA-A1/29/93/14 (15-05-1993)). Several resorts have strict policies about only purchasing legal catches.



Figure 52 Post puerulis lobster on Gan reef, 5 m, May 2009. Carapace length estimated at 1 cm

Banded coral shrimp

The banded coral shrimp (*Stenopus hispidus*) was only present in 6% of the sites – on the single site adjacent to the harbour at the Bandos Island Resort and Spa. While the shrimps are collected for the aquarium trade (Saleem, 2009) it is not considered a good indicator of human exploitation because of its scarcity, even on remote reefs.

Mollusc - Abundance and Distribution

Giant Clams – *Tridacna* spp.

Within the Indian Ocean there are eight species of giant clams, two of which are present in the Maldives, *Tridacna maxima* and *T. squamosa* (Figures 53, 54). Collectively they are locally called “gaahaka” and were the basis of a commercial export clam fishery (Ahmed, et al., 1997) before it was banned in 1993 (Notice No: FA-A1/29/93/14 (15-05-1993)). In October 2009, *Tridacna* shells (55 cm TL) were for sale in souvenir shops in Male’ for Rf12000 (US\$ = 12.75).



Figure 53 Giant clam, *Tridacna squamosa*, about 35 cm long at Vattaru, Vaavu atoll, 10 m, March 2009

The clams host a symbiotic dinoflagellate, *Symbiodinium microadriaticum*, which live freely inside the clam’s blood passages. The dinoflagellates’ products of photosynthesis assist in meeting the nutritional needs of the clams. Consequently, abundant sunlight is important for the health and rapid growth of the clam.

This is the first time *Tridacna* clams were included in the survey protocol. Consequently, there is no historical record to compare the abundance and distribution with. The survey protocol pools both species into a single taxonomic unit – clams. A total of 391 clams were recorded in 17 surveys. *Tridacna* spp. clams were found in 100% of the surveys at 3 m and 93% of the surveys at 10 m. The densities varied between atolls. The greatest numbers were observed (pooled 3 m and 10 m depth surveys representing 800 m²) at Vilingili (68 clams) Addu atoll, Hon’daafushi (43 clams) Haa Dhaalu atoll, Hithadhoo (32 clams) in Addu atoll and at Anbaraa, Vaavu atoll (33 clams) (Figure 55). The fewest clams were observed at En’boodhoofinolhu (5 clams) and Fohtheyobodufushi (6 clams), Vaavu atoll. Seventy-six percent of the clams occurred along the shallow transect (3 - 5 m) with the remaining 24% present at the deeper (10 m) transect.

Future studies should separate the species as they are different sizes at first egg release. *Tridacna maxima* first develops eggs at 8 cm TL and *T. squamosa* develops eggs at 15 cm in the Pacific Ocean (Munroe, 1993) and are probably at least five years old. Initial maturity in *Tridacna* species is known to vary with geographic location in the Pacific Ocean (SPC, 2005). Initially the clams are males and then become hermaphroditic. Very old clams generally are functionally females.



Figure 54 H. Zahir, photographing a large *Tridacna squamosa* (TL 45 cm) at Hithadhoo reef, 10 m, Addu atoll, May 2009

Setting a minimum size that corresponds to the size at which 50% of the population reaching sexual maturity is a frequently used measure that allows at least half of the giant clams to reproduce at least once before they are harvested. No data for size at initial reproduction and maximum growth length in the Indian Ocean was found. Research on the reproductive biology of *Tridacna* clams is needed to assist in the management of their stocks.

The majority (57%) of the clams were initial recruits (<10 cm T.L.) (Figure 56). Eleven percent of the juveniles occurred at Hon'daafushi. Less than 1% of the clams were >40 cm TL and are estimated to be at least 20 years old. Single individuals of this size class were only observed at Hon'daafushi, Udhafushi and Vilingili. In the Pacific Ocean, *Tridacna squamosa* and *T. maxima* are reported to grow to 40 cm and 30.5 cm total length (TL) respectively (Munro, 1993). The remainder of the clams was ranged 10 - 40 cm TL. If we assume that 50% of the clams in the 10 – 20 cm size class > 15 cm TL are reproductive, then at least 25% of the clams surveyed were likely to be females capable of releasing eggs.




Figure 55 Abundance distribution of giant clams, *Tridacna* spp., in NCRMS sites at shallow and deep transects in 2009




Figure 56 Size frequency distribution of giant clams, *Tridanca spp.*, from the 2009 NCRMS

Triton Shells

Triton is the common name given to a number of very large sea snails, predatory marine gastropods in the genus *Charonia*. Adult tritons are active predators and feed on other mollusks and starfish including adult Crown of Thorns starfish and *Culcita schmideliana*.

Many people find triton shells attractive as a design object, and so they are collected and sold as part of the international shell trade. The souvenir shops, Kaashi Boat and Gloria Maris, in Male' sell *Charonia tritonis* (TL 25cm) for Rf1300. In recent years this has doubtlessly contributed to the animals' scarcity. Traditionally Maldivians have removed the tip of the shell, or drilled a hole in the tip, and then used the shell as a trumpet. Triton shells are protected in the Maldives with an export ban imposed under fisheries regulations to stop international trade.

Although the shell was included in the survey protocol, no shells were observed on at the transects or in the roving dives.

Anthropogenic Disturbances

Over all the direct physical effects by humans on the reefs surveyed appeared to be negligible. Sand bags on the reef at En'boodhoofinolhu were quite obvious. Fishing lines were present at many sites. On occasion there appeared to be anchor damage. There were also coral observed that were damaged for no apparent reason. There was no evidence of sewage, poison fishing, blast fishing, aquarium fish collection, or invertebrate collection for the curio trade. Aquarium fish trade is an important business in the Maldives and has recently been reviewed by Saleem (2009). Only En'boodhoofinolhu has evidence of industrial activity with the increased siltation and discarded sand bags. Sand mining of the lagoon has been done manually for many decades.

Other anthropogenic disturbances such as non point source pollution are likely to affect the reefs in subtle ways. One example is the practice of fumigation on islands. This introduces toxic chemicals into the marine system without any knowledge of its effects on coral reefs, insect, bird and bat populations (Figure 57) and needs further research into the accumulation of the toxins on an island ecosystem.



Figure 57 Worker fumigating on island resort, North Male' atoll, September 2009

Sea Water Temperature Variation

Edwards, et al. (2001) used the Global sea-ice and SST data set version 2.3b (GISST2.3b) for 1° latitude x longitude areas for January 1950 to May 2000 to examine the 1998 warming anomaly in the central Maldives ($1 \pm 5^\circ$ N., $72^\circ \pm 74^\circ$ E.) without any in situ "ground truthing". The 1950 - 99 monthly SST data for the eight 1° grid squares covering the central atolls were used to calculate long-term monthly means ('normals') and standard deviations against which anomalies could be assessed. The satellite data inferred that monthly mean SST was 1.2 ± 4.0 S.D. above average during the warmest months (March – June 1998) with the greatest anomaly in May of 2.1°C (Edwards, et al., 2001). This May anomaly was 1.1°C above the highest mean monthly SST (30.3°C) expected in any 20 yr period.

Advanced Very High Resolution Radiometer (AVHRR) data has a 4.4 km spatial resolution and a seven day temporal resolution. Satellite derived SST observations only measure the temperature of a few millimeters of the ocean surface from which scientists infer temperatures several meters below the surface, where the reef community is (Quinn and Kojis, 1994). While this technique is effective for observing broad brush, large scale events it has limitations for observing localized changes, accurately assessing the temperature at which bleaching occurs, and for detecting upwellings (Quinn and Johnson, 1996; Sheppard, 2009).

NOAA satellite data is good as an early warning system to predict the general area where bleaching may occur (Liu, et al., 2005), but it is not perfect for predicting bleaching (McClanahan, et al., 2007). Monitoring programs need accurate, precise in situ underwater temperature loggers to monitor specific habitats in order to increase the resolution to better understand thermal variation. Identifying corals from similar habitats tolerate to higher temperature may help provide a population of coral that can be used for coral restoration.

In August / September / October 2009, a Reef Net Senu Ultra underwater temperature recorder was attached to a diver. The recorder was programmed to record depth and water temperatures every 10 seconds. The data was downloaded and the mean subsurface sea water temperature (S3T) was calculated based on temperatures recorded between a depth of 10 – 30 m. On 28 August 2009, the mean S3T at Akiri Channel ($4^\circ 38' 54''$ N; $73^\circ 24' 40''$ E) was 28.62°C , well within Edwards's, et al. (2001) temperature range (Figure 58). Two weeks later at the nearby site (Barakuda Point – $4^\circ 37' 49''$ N; $73^\circ 23' 37''$ E) the mean temperature was 28.50°C . However, the same week inside the atoll at Ihuru mean S3T was 28.89°C , 0.4°C warmer than channel water at Barakuda Point. Likewise, at En'boodhoofinolhu, in inner atoll site at South Male' atoll the mean S3T was 28.49°C . In late September 2009, at Han'dahaa, Gaafu Alifu atoll ($0^\circ 30' 19''$ N; $73^\circ 27' 14''$ E) the mean S3T at the same depth range was 29.55°C , over a 1°C warmer than Barakuda Point 4° latitude north. In early October at Udhafooshi, North Male' atoll, the temperature was 29.04°C , about 0.25°C above Edwards, et al. (2001) + 4 S.D. temperature range.




Figure 58 Mean satellite derived surface sea water temperature for the central atolls from 1950 – 1999 + 4 S.D. (after Edwards, 2001). Specific site data recorded between 10 - 25 m in 2009

The mean monthly S3T (MMS3T) ($n = 8698$) at 3 m at Huvafenfushi island ($4^{\circ} 22' 10.40''$ N; $73^{\circ} 22' 19.81''$ E) from August 2007 – February 2008 was 0.28°C to 1.0°C warmer than the mean monthly SST calculated from the GISST2.3b data set (Figure 59). While bleaching might be expected based on the satellite MMSST, none occurred. Corals in different habitats develop thermal tolerances allowing them to survive.

Below the surface the water is generally well mixed, probably due to a good current flow. At Rasdhoo the maximum variation in MMS3T at 8 m and 16 m is only 0.13°C , with an annual mean difference of 0.03°C (Figure 60). At these depths the MMS3T was about 0.2°C warmer than 4 S.D. above Edwards, et al., (2001) MMSST. At Barakuda Point, North Male' atoll, the temperature ranged from 29.7°C at 10 m to 29.3°C at 32 m (Figure 61) during a 43 minute period in the morning in September 2009. There is only about a 0.5°C temperature variation associated with depth even at the atoll outer reef.

The temperature logger at Rashdoo recorded several spikes in temperature in December 2008 and January 2009. The largest occurred on 11 January 2009 when the temperature dropped $>3^{\circ}\text{C}$ for a period less than eight hours (Kikinger, pers. comm.). It is uncertain what caused these upwellings and how common they are throughout the archipelago. This phenomenon needs further study with a series of underwater temperature loggers deployed throughout the country.




Figure 59 Mean S3T at coral nursery (3 m) at Huvafenfushi, North Male' atoll, from August 2007 to February 2008




Figure 60 Mean monthly sea water temperatures at 8 m and 16 m from Rasdhoo, Ari atoll, April 2008 – March 2009 compared with Edwards, et al. (2001) SST

Based on historical satellite imagery Edwards, et al. (2001) suggested that sea water temperature in the Maldives could rise of 0.16 °C per decade. They suggested that if this trend continues, by 2030 mean April SST in the central atolls will exceed, in most years, the anomaly level at which corals there are susceptible to mass bleaching. Recurrences of mass coral mortality has been predicted for several coral reef sites in the Indian Ocean including the Maldives (Sheppard, 2003).




Figure 61 Temperature by depth profile at Barakuda Point, September 2009. Y axis on the top figure is depth in meters and the °C in the bottom figure. The x axis represents dive time in minutes 10 minutes from the start of dive at 9:40am

Adaption to Sea Water Temperature Variation

Global warming may not mean death for all the coral communities around the world. Studies are showing that coral may be able to adapt to higher temperatures by forming new symbiotic associations with heat tolerant algae. In 1998, large areas of reefs were devastated throughout the world. Reefs surviving in Kenya, Panama and the Arabian Gulf provided clues that some zooxanthellae were heat tolerant. Field studies showed that coral in these areas contained more D-type algae than non heat tolerant C-type algae (Baker, et al., 2004). This was also demonstrated by experimental studies in Panama, Guam and Great

Barrier Reef. The researchers showed that the common Indo-Pacific coral *Acropora millepora* was able to increase its upper thermal limit by 1 °C - 1.5 °C when it changed from C-type to D-type algae.

The genus *Symbiodinium* is diverse, and many corals are relatively flexible in the type(s) of algal symbiont they contain, although one type is usually dominant in any given species and environment. The conditions needed for a coral to change its algal symbionts are unknown. Baker et al. (2004) considered that the symbiont changes are a common feature of severe bleaching and mortality events, and predicted that these adaptive shifts will increase the resistance of these recovering reefs to future bleaching. The implications are that if this mechanism of acclimatization is as widespread as it appears, coral reefs may have significantly more flexibility to respond to climate change than previously thought. However, it is not known whether symbiont change alone is sufficient for coral reefs to adapt to current climate warming predictions of 1-3 °C. Even if this acclimatization mechanism can match global warming, the structure of the world's coral reefs may change dramatically with fewer species and probably lower coral cover (Berkelmans and Oliver, 1999).

Future Coral Reef Monitoring Efforts and Awareness Development

Assessing the recovery of the reefs in the Maldives is like the story of three blindmen each describing an elephant by touching the trunk, leg or tail. There is considerable variation among reefs and among habitats within reefs. This variation is a function of the wide latitudinal range of the atolls, the structure of the atolls, and chance. These factors create considerable variation in the abiotic factors that influence reef development and, therefore, considerable variation among reefs.

While this survey has expanded the depth at which reef transect data was collected (10 m), future surveys should include sites at depths of 15 – 20 m and include the outer atoll reef crest and slope habitats (Figure 62). Expansion of sampling sites to deeper depths and more exposed sites would require an experienced dive team with advanced diver qualifications.




Figure 62 South Male' atoll, outer reef, 15 m, August 2009. Future surveys need to include coral communities on the outer reef slope of the atolls. Lateral visibility estimated at over 35 m

The recovery of most of the reefs surveyed by MRC during the past decade is remarkable and suggests the question of where were the new recruits coming from. Zahir (2006) observed that recruitment was not a limiting factor for reef recovery. The currents in the Maldives are very unpredictable. On each dive a dive master enters the water to assess the direction and strength of the current which seems to move independently of tidal phase and wind direction. A detailed study of currents around the Maldives would help to determine the larval dispersal for both invertebrate and reef fish stocks. This information is necessary to better understand and manage these larval source communities.

One needs to be careful in extrapolating predictions to a larger scale based on specific observations to a single reef. Schuhmacher, et al. (2006) observed an "... ongoing deterioration of the reef or at least a prolonged time for recovery..." for the Komandoo reef flat based on seven years of study. Even the results of a spatially diverse data set based on a 150 m sample of a reef can be misleading. For example, Zahir (2006) reported a change in percent coral cover in the 16 NCRMS sites from -65% to 5916%. Because of the very low level of coral cover small changes in the presence of coral result in large percent changes. The coefficient of variation was greater than 70% for 12 of the 15 sites in 1999 and 9 of 15 sites in 2005. Ten of the 15 sites had <9% coral cover and only one site (Fesdhoo) had moderate coral cover (>30%). Coral cover in the early years of the NCRMS was sparse and variable both within a reef at a given depth and throughout the country. Consequently, small changes in cover could result in misleading inferences if extrapolated to larger dimensions.

For centuries the coral reefs have adapted to changes in sea levels associated with climate change. Kench, et al. (2009b) has demonstrated by drilling through several reefs in South Maalhosmadulu atoll that healthy coral reefs produced sufficient skeletal material to allow reef islands to accommodate rising sea levels on the order that is projected for the next century. The question today is: Are the reefs sufficiently healthy to continue to keep up with sea level rise and maintain the islands or are global and local stresses to the reefs so intense as to diminish the growth and reproduction of the corals resulting in submergence of all or some of the islands of the Maldives?

Uninhabited or lightly inhabited islands may survive. Sand can accumulate and raise the height of the islands. Beach rock can form and storms can carry coral rocks about sea level. They can't do it on the heavily inhabited, manufactured islands without destroying the infrastructure. Male' will need to be manually protected even if reefs keep up as the sand produced will just be washed into the lagoon. As sea level raises the freshwater lens in Male' and other islands will become increasingly saline. Roof top water catchments for storage in underground cisterns may be a partial solution. Other islands that have already physically destroyed parts of their reef and are creating an environment that is probably slowing coral growth, reproduction, and recruitment are also in need of survival strategies.

Future monitoring efforts should focus on providing a scientific framework for the management and justification of Marine Protected Areas (MPAs). Marine Protected Areas in the Maldives lack the necessary elements for effective management (Mohamed, 2007). She considered these MPAs to be only "paper parks" in that they are protected through legislation with no real funding for outreach and education and enforcement to protect them. She also suggested that the main reason for this the lack of enforcement was little awareness and understanding of the true value of reef resources.

The only NCRMS site within a MPA was Hithadhoo, Addu atoll. Hithadhoo reef had been included in a resource assessment for the establishment of the Maldives Protected Area System (Ministry of Environment, Energy and Water, 2006), prior to its recent inclusion in the NCRMS. The Vattaru site was within a kilometer of the Vattaru MPA. Future coral reef monitoring efforts should include MPAs in order to assess their effectiveness. Without proper monitoring it is difficult to assess the effectiveness of these MPAs in protecting the reefs. The NCRMS could support the MPA concept by providing information within an MPA and at a suitable control site.

Socio economic studies should also be undertaken to assess the usage of the parks by water sports organizations and the effectiveness of the parks in enhancing nearby fish stocks. Coral reef monitoring efforts need to be linked with the education curriculum and with social organizations in each atoll to foster greater adult understanding of science based management principals. More effort could be made to develop swimming, diving and snorkeling skills in Maldivian youth on all islands. The many Maldivians that NJQ has shared dives with are among the most passionate conservationists he has met.

The threat of increasing atmospheric CO₂ on coral reef calcification poses a chronic and increasing threat to coral reefs. However, it is also a creeping and nearly invisible stress that receives much less attention than the more visible and acute coral reef problems like bleaching and diseases. Long term time series

data for abiotic parameters need to be included in a NCRMS program. Data loggers for subsurface seawater temperature are inexpensive and reliable and should be deployed at all survey sites. Other recorders for salinity, pH, dissolved oxygen, wave height and nutrient levels require more maintenance and could be deployed during surveys and at an easily accessible site near Male' where they can be frequently monitored and maintained.




Figure 63 Photosynthesis is a process that converts carbon dioxide into organic compounds, especially sugars, using the energy from sunlight. In the sea, instead of a tomato plant, photosynthesis is done by plankton, coral, algae and sea grass

Incident waves and their interaction with coral reef platforms are considered the main mechanism controlling the formation and stability of low lying coral reef islands. Large differences in incident wave energy between the SW and NE monsoons can affect the pattern of beach erosion on islands (Kench, et al., 2009a). Wind driven incident waves are both refracted and diffracted around the reef platform creating currents which result in nodal location for sediment deposition (Kench, et al., 2009a). The seasonal occurrence of various intensities and the direction of wind results in movement of islands. While scientists monitor changes in island configuration, few place the movement within a meteorological or diffraction/refraction context. Consequently, when beaches erode on inhabited islands, climate change is considered the cause, rather than considering natural variations in wind intensity and direction or man-made changes in current patterns and wave action caused by the placement of breakwaters and structures or dredging of the lagoon. The placement of automated meteorological stations on several islands within an atoll would help provide better data for understanding localized climatic conditions which influence incident waves and the resultant island movement.

It is hypothesized that the Maldives is a CO₂ sink and could actually sell carbon credits on the international market. This of course requires a detailed scientific survey. The implementation of a series of underwater fluorometers would begin to measure the photosynthesis of corals, algae, plankton and sea grasses. Photosynthesis is a process that converts carbon dioxide into organic compounds, especially sugars, using the energy from sunlight (Figure 63). Photosynthesis is vital for life on Earth. As well as maintaining the normal level of oxygen in the atmosphere, nearly all life either depends on it directly as a source of energy, or indirectly as the ultimate source of the energy in their food.

The Government of the Republic of Maldives has applied for financing from the International Development Association (IDA) to cover the cost of the Maldives Environmental Management Project. The project seeks to:

- 1) strengthen Maldives' existing national coral reef monitoring program by the inclusion of additional monitoring stations and parameters such as water quality and proxies for reef health,
- 2) support the development of a community-based coral reef monitoring and community awareness program aimed at improving the understanding about the importance of coral reef systems to the integrity of the island and atoll ecosystem. A report is currently under review and the acceptance of the final report will determine the scope of future coral reef monitoring activities in the Maldives.

References

- Adam, M.S., Anderson, R.C. and Shakeel, H. 1997. Commercial exploitation of reef resources: examples of sustainable and non sustainable utilization from the Maldives. in: Proceedings of the 8th International Coral Reef Symposium. 2:2015-2020.
- Ahmed, H., Mohamed, S. and Saleem, M.R. 1977. Paper 4. Exploitation of reef resources: Beche-de-mer, Reef Sharks, Giant Clams, Lobsters and Others. in: Nickerson, D.J. and Maniku, M.H. (eds.) Workshop on integrated reef resource management in the Maldives. Bay of Bengal Programme. pp. 93-116.
- Allison, W.R. 1995. Changes in the Maldivian reef system. *Coastal Management in Tropical Asia* 4:6-8.
- Allison, W.R. 1996. Snorkeler damage to reef corals in the Maldivian Islands. *Coral Reefs* 15:215-218.
- Allison, W. 1999. Maldives/GCRMN pilot reef monitoring study: Post-bleaching reef status after the mass bleaching in 1998. Marine Research Centre, Ministry of Fisheries, Agriculture and Marine Resources, Maldives.
- Anderson, R.C. 2006. Baitfish and reef fish analysis and management. FAO Report. pp. 55.
- Arthur, R., Done, T.J., Marsh, H. and Harriott, V. 2006. Local processes strongly influence post-bleaching benthic recovery in the Lakshadweep Islands. *Coral Reefs* 25:427-440.
- Aubert, O. and Drozler, A. 1992. General Cenozoic evolution of the Maldives carbonate system (Equatorial Indian Ocean): *Bulletin des Centres de Recherches Exploration Production Elf Aquitaine* 16:113-136.
- AusAID, 2005. An assessment of damage to Maldivian coral reefs and baitfish populations from the Indian Ocean tsunami. Australian Government / Government of the Republic of the Maldives report. pp. 68.
- Baker, A.C., Craig, J., Starger, C.J., McClanahan, T.R. and Glynn, P.W. 2004. Corals' adaptive response to climate change. *Nature* 430:791.
- Berkelmans, R. and Oliver, J. K. 1999. Large-scale bleaching of corals on the Great Barrier Reef. *Coral Reefs* 18: 55-60.
- Bianchi, C.N., Morri, C., Pichon, M., Benzoni, F., Colantoni, P., Baldelli, G. and Sandrini, M. 2006. Dynamics and pattern of coral recolonization following the 1998 bleaching event in the reefs of the Maldives. *Proceedings of the 10th International Coral Reef Symposium* 1:30-37.
- Brown B.E. and Dunne, R.P. 1988. The impact of coral mining on coral reefs in the Maldives. *Environmental Conservation* 15:159-165.
- Burt, J., Bartholomew, A. and Usseglio, P. 2008. Recovery of corals a decade after a bleaching event in Dubai, United Arab Emirates. *Marine Biology* 154:27-36.
- Carpenter, K.E. et al. 2008. One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science* 321:560-563.
- Chen, C.A. and Dai, C.F. 2004. Local phase shift from *Acropora*-dominant to *Condylactis*-dominant community in the Tiao-Shi Reef, Kenting National Park, southern Taiwan. *Coral Reefs* 23:508.
- Ciarapica G. and Passeri L. 1993. An overview of the Maldivian coral reefs in Felidu and North Malé atolls (Indian Ocean): Platform drowning by ecological crises. *Facies* 28:3-66.
- Clark, S., Akester, S. and Naeem, H. 1999. Status of the coral reef communities in North Malé Atoll, Mal-

- dives: Recovery following a severe bleaching event. Report to the Ministry of Home Affairs, Housing and Environment, February 1999. pp. 13.
- Clark, S. 2000. Evaluation of succession and coral recruitment in Maldives. in: Coral Reef Degradation in the Indian Ocean. Souter, D., Obura D. and Lindén, O. (eds.), Stockholm: Cordio, 2000. pp. 169-175.
- Clua, E., Legendre, P., Vigliola, L., Magron, F., Kulbicki, M., Sarramegna, S., Labrosse, P. and Galzin, R. 2006. Medium scale approach (MSA) for improved assessment of coral reef fish habitat. *J Exp Mar Biol Ecol* 333:219–230.
- Conand, C. 1990. The fishery resources of Pacific island countries Part 2. Holothurians. FAO Technical Paper 272.2.
- Conand, C. 1990. The fishery resources of Pacific island countries Part 2. Holothurians. FAO Technical Paper 272.2.
- Edwards, A.J., Clark, S., Zahir, H., Rajasuriya, A., Naseer, A. and Rubens, J. 2001. Coral bleaching and mortality on artificial and natural reefs in Maldives in 1998, sea surface temperature anomalies and initial recovery. *Marine Pollution Bulletin* 42:7-15.
- English, S., Wilkinson, C. and Baker, V. (eds). 1994. Survey Manual for Tropical Marine Resources. ASEAN-Australian Marine Science Project: Living Coastal Resources. Australian Institute of Marine Science, Townsville. pp. 368.
- Gillett, R. 2004. Aspects of fisheries management in the Maldives. FAO/FishCode Review No.2: 1-54.
- Goreau, T., McClanahan, T.R., Hayes, R. and Strong, A. 2000. Conservation of coral reefs after the 1998 coral bleaching event. *Conservation Biology* 14:5-15.
- Graham, N.A.J., Wilson, S.K., Jennings, S., Polunin, N.V.C., Bijoux, J.P. and Robinson, J. 2006. Dynamic fragility of oceanic coral reef ecosystems. *Proceedings of the National Academy of Sciences*. 103:8425-8429.
- Hass, H. 1961. Expedition in Unbekannte; Ein Bericht iiber die expedition des Forschungsschiffe "Xarifa" zu den Malediven und Nikobaren. Berlin, pp. 167.
- Heyward, A.J. and Negri, A.P. 1999. Natural inducers for coral larval metamorphosis. *Coral Reefs* 18:273–279.
- Hodgson, G. 1999. A global coral reef assessment of human effects on coral reefs. *Marine Pollution Bulletin* 38:345-355.
- Hodgson, G. and Liebler, J. 2002. The global coral reef crisis trends and solutions. *Reef Check*. pp. 77.
- Hodgson, G., Hill, J., Kiene, W., Maun, L., Mihaly, J., Liebler, J., Shuman, C. and Torres, R. 2006. Reef Check Instruction Manual: A Guide to Reef Check Coral Reef Monitoring. Reef Check Foundation, Pacific Palisades, California, USA.
- Hughes, T.P., Reed, D.C. and Boyle, M.J. 1987. Herbivory on coral reefs: Community structure following mass mortalities of sea urchins. *J. Exp. Mar Biol. Ecol.* 113:39-59.
- Hughes, T.P. 1994. Catastrophes, phase shifts, and large scale degradation of a Caribbean coral reef. *Science* 265:2547-1552.
- Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L., Moltschanowskyj, N., Pratchett, M.S., Steneck, R.S. and Willis, B. 2007. Phase shifts, herbivory, and the resilience of coral reefs to climate change. *Current Biology* 17(4):360–365.
- Jonklass, R. 1961. A preliminary investigation of the spiny lobster resources in the Maldivian Islands. Unpublished report. pp. 7.

- Joseph, L. 1992. Review of the beche-de-mer (sea cucumber) fishery in the Maldives. BOBP/WP/79: 1-31.
- Kench, P.S., McLean, R.F. and Nichol, S.L. 2005. New model of reef-island evolution: Maldives, Indian Ocean. *Geology* 33(2):145-148.
- Kench, P.S., Brander, R.W., Parnell, K.E. and McLean, R.F. 2006. Wave energy gradients across a Maldivian Atoll: implications for island geomorphology. *Geomorphology* 81:1-17.
- Kench, P.S., Mclean, R.F., Brander, R.W., Nichol, S.W., Smithers, S.G., Ford, M.R., Parnell, K.E. and Aslam, M. 2006. Geological effects of tsunami on mid-ocean atoll islands: The Maldives before and after the Sumatran tsunami. *Geology* 34(3):177-190.
- Kench, P.S., Brander, R.W., Parnell, K.E. and O'Callaghan, J.M. 2009a. Seasonal variations in wave characteristics around a coral reef island, South Maalhosmadulu atoll, Maldives. *Marine Geology* 262:116-129.
- Kench, P.S., Smithers, S.G., McLean, R.F. and Nichol, S.L. 2009b. Holocene reef growth in the Maldives: Evidence of a mid-Holocene sea-level highstand in the central Indian Ocean. *Geology* 37:455-458.
- Kojis, B.L. and Quinn, N.J. 2001. The importance of regional differences in hard coral recruitment rates for determining the need for coral restoration. *Bulletin of Marine Science* 69(2): 967-974.
- Lasagna, R., Albertelli, G., Giovannetti, E., Grondona, M., Milani, A., Morri, C. and Bianchi, C.N. 2008. Status of Maldivian reefs eight years after the 1998 coral mass mortality. *Chemistry and Ecology* 24:67-72.
- Lasagna, R., Albertelli, G., Colantoni, P., Morri, C. and Bianchi, C.N. 2009. Ecological stages of Maldivian reefs after the coral mass mortality of 1998. *Facies* 1-11.
- Ledlie, M.H., Graham, N.A.J., Bythell, J.C., Wilson, S.K., Jennings, S., Polunin, N.V.C. and Hardcastle, J. 2007. Phase shifts and the role of herbivory in the resilience of coral reefs. *Coral Reefs* 26:641-653.
- Littler, M.M. and Littler, D.S. 1997. Disease-induced mass mortality of crustose coralline algae on coral reefs provides for the conservation of herbivorous fish stocks. *Proceedings of the 8th International Coral Reef Symposium* 1:719-724.
- Loch, K., Loch, W., Schuhmacher, H. and See, W.R. 2002. Coral recruitment and regeneration on a Maldivian reef 21 months after the coral bleaching event of 1998. *Marine Ecology* 23(3): 219-236.
- Lovatelli, A., Conand, C., Purcell, S., Uthicke, S., Hamel, J.F., Mercier, A. 2004. Advances in sea cucumber aquaculture and management. *FAO Fisheries Technical Paper-T463*.
- Lui, G., Strong, A.E., Skiving, W., Arzayus, L.F. 2005. Overview of NOAA coral reef watch program's near-real time satellite global coral bleaching monitoring activities. *Proceedings 10th International Coral Reef Symposium* 1:1783-1793.
- Massin, C., Rasolofonirina, R., Conand, C. and Samyn, Y. 1999. A new species of *Bohadschia* (Echinodermata: Holothuroidea) from the Western Indian Ocean with a redescription of *Bohadschia subrubra* (Quoy & Gaimard, 1833). *Bull. Inst. Roy. Sci. Nat. Belg.* 69: 151-160.
- McClanahan, T.R. 2000. Bleaching damage and recovery potential of Maldivian coral reefs. *Marine Pollution Bulletin* 40:587-597.
- McClanahan, T.R., Ateweberhan, M., Ruiz Sebastian, C., Graham, N.A.J., Wilson, S.K., Bruggemann, J.H. and Guillaume, M.M.M. 2007. Predictability of coral bleaching from synoptic satellite and in situ temperature observations. *Coral Reefs* 26:695-701.
- McCook, L.J., Jompa, J., Daiz-Pulido, G. 2001. Competition between corals and algae on coral reefs: a review of evidence and mechanisms. *Coral Reefs* 19:400-417.

- Ministry of Environment, Energy and Water. 2006. Assessment of Eidhigali Kulhi and Koatthey Area, S. Hithadhoo, Maldives. Male', Maldives. pp. 137.
- Ministry of Planning and National Development, 2008. Official Atlas of the Maldives, Novelty Printers, Male', Maldives. pp. 61
- Mohamed, M. 2007. Economic valuation of Coral Reefs: A Case Study on the Costs and Benefits of Improved Management of Dhigali Haa, a Protected Area in Baa Atoll, Maldives. M.Sc. thesis, University of Canterbury. pp.163.
- Munro, J. 1993. Giant clams. in: Wright, A. and Hill, L. (eds). Nearshore Marine Resources of the South Pacific: Information for Fisheries Development and Management. pp. 431-449. Forum Fisheries Agency, Honiara, Solomon Islands.
- Muthiga., N. 2008. Field observations of Sea Cucumbers at North Male Atoll in the Maldives. SPC Beche-de-Mer Information Bulletin. 27: p33-37
- Naeem, I., Rasheed, A., Zuhair, M. and Riyaz, M. 1998. Coral bleaching in the Maldives –1998. Survey carried out in the North and South Malé atolls. pp. 14.
- Pacer-Rosario, G. 1987. The effect of substratum on the growth of *Terpios*, an encrusting sponge which kills coral. *Coral Reefs* 5:192-200.
- Pichon M. and Benzoni, F. 2007. Taxonomic re-appraisal of zooxanthellate Scleractinian corals in the Maldives Archipelago. *Zootaxa* 1441: 21–33.
- Price, A.R.G. , Harris, A., McGowan, A., Venkatachalam A.J., Sheppard C.R.C. 2009. Chagos feels the pinch: assessment of holothurian (sea cucumber) abundance illegal harvesting and conservation prospects in British Indian Ocean Territory. *Aquatic Conservation: Marine and Freshwater Ecosystems*. DOI: 10.1002/aqc.1054
- Quinn, N.J. and Kojis, B.L. 1994. Evaluation of the use of AVHRR satellite imagery and in situ obtained subsurface sea water temperatures for monitoring coastal marine communities in the Caribbean Sea. *Proc 2nd Thematic Conf on Remote Sensing for Marine & Coastal Environments*. New Orleans, 1:653-664.
- Quinn, N.J. and Johnson, D.W. 1996. Cold water up welling cover Gulf of Oman coral reefs. *Coral Reefs* 15:218.
- Rajasuriya, A. and Karunaratna, C. 2000. Post-bleaching status of the coral reefs of Sri Lanka. in: *Coral Reef Degradation in the Indian Ocean*. Souter, D., Obura, D. and Lindén, O. (eds.), Stockholm: Cordio, 2000. pp. 54–63.
- Reichenbach, N. 1999. Ecology and fishery biology of *Holothuria fuscogilva* (Echinodermata: Holothuroidea) in the Maldives, Indian Ocean. *Bulletin of Marine Science*, 64(1): 103-111.
- Riyaz, M., Shareef, M. and Elder, D. 1998. Coral bleaching event: Republic of Maldives, May 1998. Ministry of Home Affairs, Housing and the Environment.
- Rogers, C.S. and Miller, J. 2006. Permanent 'phase shifts' or reversible declines in coral cover? Lack of recovery of two coral reefs in St. John, US Virgin Islands, *Marine Ecological Progress Series* 306:103–114.
- Saleem, M. 2009. The aquarium fish trade in the Maldives. *Proceedings of the 11th International Coral Reef Symposium*. (in press).
- Sattar, S.A., 2008. Review of the Reef Fishery of Maldives. Marine Research Centre, Male', Republic of Maldives. 62pp.
- Sattar, S.A. and Adam, M.S, 2005. Review of Grouper Fishery of the Maldives with Additional Notes on the

- Faafu Atoll Fishery. Marine Research Centre, Male', Republic of Maldives. 54pp.
- Scheer, G. 1971. Coral reefs and coral genera in the Red Sea and Indian Ocean. Symposium of Zoological Society of London. 28:329-367.
- Scheer, G. 1972. Investigation of coral reefs in the Maldivian Islands with notes on lagoon patch reefs and the method of coral sociology. in: Proceedings of the 1st International Coral Reef Symposium. pp. 87-120.
- Scheer, G. 1974. Investigation of coral reefs at Rasdu Atoll in the Maldives with the quadrat method according to phytosociology. in: Proceedings of the 2nd International Coral Reef Symposium. 2:655-670.
- Schuhmacher, H., Loch, W. and Loch, K. 2002. Post-bleaching growth reveals *Diploastrea helianthus* to be a coral methusalum. Coral Reefs. 21:344-345.
- Schuhmacher, H., Loch, K., Loch, W. and See, W.R. 2005. The aftermath of coral bleaching on a Maldivian reef—a quantitative study. Facies 51:80–92.
- Schuhmacher, H., Loch, K., Loch, W. and See, W.R. 2006. Post-bleaching development of a Northern Maldivian Reef. Proceedings of the 10th International Coral Reef Symposium. 657-663.
- Sheppard, C.R.C. 1999. Coral decline and weather patterns over 20 years in the Chagos Archipelago, central Indian Ocean. *Ambio* 28:472-478.
- Sheppard, C.R.C. 2003. Predicted recurrences of mass coral mortality in the Indian Ocean. *Nature* 425:294-297.
- Sheppard, C.R.C. 2009. Large temperature plunges recorded by data loggers at different depths on an Indian Ocean atoll: comparison with satellite data and relevance to coral refuges. *Coral Reefs* 28:399-403.
- Sheppard, C.R.C. and Wells, S. 1988. *Coral Reefs of the World*. Gland, Switzerland, UNEP/IUCN. pp. 389.
- Sheppard, C.R.C., Spalding, M., Bradshaw, C. and Wilson, S. 2002. Erosion vs. recovery of coral reefs after 1998 El Niño: Chagos reefs, Indian Ocean. *Ambio* 31:40–48.
- Sheppard, C.R.C., Dixon, D.J., Gourlay, M., Sheppard, A. and Payet, R. 2005. Coral mortality increases wave energy reaching shores protected by reef flats: examples from the Seychelles. *Estuarine and Coastal Shelf Science*. 64:223–234.
- Sheppard C.R.C., Harris A. and Sheppard, A. 2008. Archipelago-wide coral recovery patterns since 1998 in the Chagos Archipelago, Central Indian Ocean. *Mar. Ecol. Prog. Ser.* 362:109–117.
- Sluka, R. and Miller, M.W. 1998. Coral mining in the Maldives. *Coral Reefs* 17:288.
- Smith, L. D., Gilmour, J.P. and A. J. Heyward, A.J. 2008. Resilience of coral communities on an isolated system of reefs following catastrophic mass-bleaching. *Coral Reefs* 27:197–205.
- Spalding, M.D. and Jarvis, G.E. 2002. The impact of the 1998 coral mortality on reef fish communities in the Seychelles. *Marine Pollution Bulletin* 44:309-321.
- SPC. 2005. Size limits and other coastal fisheries regulations used in Pacific islands region. Coastal fisheries management and fisheries information sections of the Secretariat of Pacific Community. pp. 32.
- Spencer-Davies, P., Stoddart, D.R. and Sigeo, D.C. 1971. Reef forms of Addu Atoll, Maldivian Islands. *Symp. Zool. Soc. Lond.* 28:217–259.
- Solandt, J.-L. and Wood, C. 2008. Maldives Reef Survey – June 13 – 30 2008. Marine Conservation Society. pp. 18.

- Stoddart, D.R., Spencer-Davies, P. and Keith, A.C. 1966. Geomorphology of Addu Atoll. *Atoll Research Bulletin* 116:13–41.
- United Nations Environmental Programme. 2003. <http://coral.unep.ch/atlaspr.htm>
- Veron, J.E.N., et al. 2009. The coral reef crisis: The critical importance of <350 ppm CO₂. *Marine Pollution Bulletin* 58: 1428-1436.
- Wallace C.C. and Zahir, H. 2007. The “Xarifa” expedition and the atolls of the Maldives, 50 years on. *Coral Reefs* 26:3–5.
- Wapnick, C., Precht W.F. and Aronson, R.B. 2004. Millennial-scale dynamics of staghorn coral in Discovery Bay, Jamaica. *Ecology Letters* 7: 354-361
- Wilkinson, C. 1998. The 1997-1998 mass bleaching event around the world. in: Wilkinson, C.R. (ed.) *Status of coral reefs of the world 1998*. Australian Institute of Marine Science, Townsville. pp. 15-38.
- Wilson, S.K., Graham, N.A.J. and Polunin, N.V.C. 2007. Appraisal of visual assessments of habitat complexity and benthic composition on coral reefs. *Marine Biology* 151:1069–1076.
- Wood, E. 1987. Recent reef degradation (bleaching). In: Engineering Geology Ltd and Tropical Management Consultants Ltd, *Geological, Technical and ecological studies of selected atolls of the Republic of Maldives*. Report No. 426/FE/1187B. Department of Public Works and Labor, Maldives, pp. 35-44.
- Woodley, J.D. 1989. The effects of Hurricane Gilbert on coral reefs at Discovery Bay. in: Bacon, P. (ed.) *Assessment of the Economic Impacts of Hurricane Gilbert on Coastal and Marine Resources in Jamaica*. pp 79-82. CEP Technical Report No. 4, UNEP Caribbean Environment Programme, Kingston, Jamaica. pp. 87.
- Wright, A. 1992. The Maldives fishery resources assessment and requirements for development and management. in: *Maldives fishery sector strategy study. Final report Vol 2*. Primex-GOPA-TPC. pp. 95.
- Zahir, H. 2000. Status of the coral reefs of Maldives after the bleaching event in 1998. in: *Coral Reef Degradation in the Indian Ocean*. Souter, D., Obura, D. and Lindén, O. (eds.), Stockholm: Cordio, 2000. pp. 64–68.
- Zahir, H. 2002a. Status of the coral reefs of Maldives. in: *Coral Reef Degradation in the Indian Ocean*, Lindén, O., Souter, D., Wilhelmsson, D. and Obura, D. (eds.), Stockholm: Cordio, 2002. pp. 119–124.
- Zahir, H. 2002b. Assessing bioerosion and its effect on reef structure following a bleaching event in the Maldives. in: *Coral Reef Degradation in the Indian Ocean*, Lindén, O., Souter, D., Wilhelmsson, D. and Obura, D. (eds.), Stockholm: Cordio, 2002. pp. 135–138.
- Zahir, H. 2006. Patterns of coral community recovery in the Maldives following mass bleaching in 1998. MPhil. thesis, Newcastle University, pp. 118.
- Zahir, H., Naeem, I., Rasheed, A. and Haleem, I. 1998. *Reef Check Maldives: Reef Check 1997 and 1998*. Marine Research Section, Ministry of Fisheries, Agriculture and Marine Resources, Male’, Republic of Maldives.
- Zahir, H., Clark, S., Ajla, R. and Saleem, M. 2002. Spatial and temporal patterns of coral recruitment following a severe bleaching event in the Maldives. in: *Coral Reef Degradation in the Indian Ocean*, Lindén, O., Souter, D., Wilhelmsson, D. and Obura, D. (eds.), Stockholm: Cordio, 2002. pp. 125-134.
- Zahir, H., Allison, W., Dews, G., Gunn, J., Rajasuriya, A., Solandt, J.-L., Sweatman, H., Tamelander, J., Thompson, A. and Wakeford, M. 2006. Post-Tsunami Status of the Coral Reefs of the Islands and Atolls of the Maldives. in: Wilkinson, C.R., Souter, D. and Goldberg, J. (eds.) *Status of coral reefs in tsunami affected countries: 2005*. Australian Institute of Marine Science. Townsville. pp. 111-120.



Marine Research Centre
H. White Waves, Moonlight Higun, 20025, Malé, Republic of Maldives